## Prevalence of Cardiovascular Disease and Associated Risk Factors among Adult Population Attending Fitness Center/Gymnasium in Arar

#### Anshoo Agarwal<sup>1</sup>\*, Rahma Hamayun Fazeel<sup>2</sup>, Hana El-Sayed Abdel Rahman<sup>3</sup>, Atheer Muteb K Alenezi<sup>1</sup>, Dhari Atallah S Alshammari<sup>1</sup>, Manal Hajea E Alenzi<sup>1</sup>, Talal Ahmed Albalawi<sup>1</sup>, Yazed Muteb K Alenezi<sup>4</sup>, Rahaf Hajea E Alenazi<sup>4</sup>

<sup>1</sup>Department of Pathology, Faculty of Medicine, Northern Border University, Arar, Saudi Arabia. <sup>2</sup>Department of Pharmacology, Faculty of Medicine, Northern Border University, Arar, Saudi Arabia. <sup>3</sup>Department of Family and Community Medicine, Faculty of Medicine, Northern Border University, Arar, Saudi Arabia. <sup>4</sup>Faculty of Pharmacy Northern Border University, Rafha, Saudi Arabia.

#### Abstract

Cardiovascular disease is the most frequent cause of death in Saudi Arabia; next to road traffic accidents. Both resistance and aerobic exercises provide the greatest benefits as the best exercise modality for the prevention of cardiovascular diseases. To determine the prevalence of cardiovascular disease and associated risk factors among the adult population attending the Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia. In a randomized cross-sectional prospective study, unselected adult population attending a Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia was screened for Obesity, Diabetes mellitus, and hypertension and its risk factors. The outcomes of this study were analyzed using the SPSS program using a pre-tested questionnaire. The study included 213 participants more than half of them 57.7% 20-30 years old, 63.8% females, and 36.2% males. 72.3% of participants joined the club for general health and fitness, 33.8% to lose weight, and 22.1% to spend spare time. Only 10.8% had been diagnosed with high blood pressure. 89.2% correctly answered that practicing physical activity can reduce the risk of cardiovascular disease. 84% said that walking can prevent the risk of cardiovascular disease and 68.5% said that usual physical activity for 150 minutes a week will reduce your chance of having a heart attack or stroke. Participants showed good awareness of cardiovascular disease risk factors among participants which was in the same line with other reported studies. Also, they know the role of physical activity in reducing the risk of cardiovascular disease risk.

Keywords: Fitness, CVD, Cardiovascular risk, Gym, Fitness center, KSA

### **INTRODUCTION**

Cardiovascular diseases have become a global epidemic and are an important public health challenge in both economically developing and developed countries [1]. The global prevalence of these risk factors is being increased every day. Cardiovascular disease is the frequent cause of death in Saudi Arabia; next to road traffic accidents. Hypertension that is not controlled is related with an end-organ damage which includes heart disease, stroke, blindness, and renal disease [2]. One of the most damaging complications of hypertension may be prevented with proper blood pressure control [3]. As there is a high prevalence of cardiovascular risk factors in the Kingdom of Saudi Arabia and these days many people are aware of it but many are still ignorant. We wish to study how many adults in Arar in the Kingdom of Saudi Arabia join a gymnasium and fitness centre to do regular exercise to reduce the risk for cardiovascular diseases. Many of these adults attending fitness centres may have been spending most of the day at home and therefore are at risk of various metabolic disorders due to ignorance about the food they eat that may be salt-laden or high in oil adding to their physically inactive lifestyle [4]. This subpopulation requires intensive care comprising intensive health education in cardiovascular risk

factors management [5, 6]. Massive population screening for cardiovascular risk is needed and adequate blood pressure and glycemic control are essential to reduce associated mortality and morbidity [7].

Aerobic exercise training yields important health benefits. Just 15 minutes of moderate-intensity exercise a day can reduce the risk of premature death from cardiovascular disease [8, 9]. The Health benefits of exercise are usually

Address for correspondence: Anshoo Agarwal, Department of Pathology, Faculty of Medicine, Northern Border University, Arar, Saudi Arabia. dranshoo30@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to cite this article: Agarwal A, Fazeel RH, Abdel Rahman HES, Alenezi AMK, Alshammari DAS, Alenzi MHE, et al. Prevalence of Cardiovascular Disease and Associated Risk Factors among Adult Population Attending Fitness Center/Gymnasium in Arar. Arch Pharm Pract. 2022;13(3):127-33. https://doi.org/10.51847/SBqqCZtmzv

based on endurance training, such as running and cycling [8]. Resistance training and strength training are already known to reduce the risk of type 2 diabetes and improve bone health, but much research is lacking on its impact on the development of cardiovascular disease and metabolic syndrome. Not done [9]. People who follow exercise guidelines (two or more sessions per week) have a lower risk of developing cardiovascular diseases [10-12]. Resistance exercise for even less than one hour per week is associated with a 29 percent lower risk [13-15]. Both resistance and aerobic exercises provide the greatest benefits as the best exercise modality for the prevention of cardiovascular diseases [16].

We aim to study if our study results can indicate that exercise, such as two 30-minute sessions per week, has the most beneficial effect among adults attending a gymnasium. Fitness center. Based on our results we wish to create awareness by giving medical recommendations for preventing metabolic syndrome and future cardiovascular disease."

### Aims

To determine the prevalence of cardiovascular disease and associated risk factors among the adult population attending the Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia.

### **Objectives**

- 1. To study the prevalence and awareness of obesity among the adult population attending a Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia.
- 2. To study the prevalence and awareness of cardiovascular disease risk factors among the adult population attending the Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia.
- 3. To study the effect of different types of exercises undertaken by the adult population attending the Fitness center/Gymnasium in reducing cardiovascular risk factors.

## MATERIALS AND METHODS

In a randomized cross-sectional prospective study, unselected adult population attending a Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia was screened for Obesity, Diabetes mellitus, and hypertension and its risk factors. To study the demographic details and to get data for our research objectives, a structured questionnaire was given randomly to the participants who were taken from most of the Fitness centers/Gymnasium for adults in Arar, Kingdom of Saudi Arabia.

Step 1: The questionnaire addressing personal information, bio-data, habits, and practices related to the risk factors of cardiovascular diseases as well as their educational status [includes age, nationality, occupation, education, employment status, marital status, diet intake, fast food intake, smoking, drinking energy drinks, hypertension awareness, history of diabetes or any other cardiovascular disease, the status of exercise].

Stage 2: Measurement of blood pressure by sphygmomanometer for three consecutive days with one-week intervals for more accurate assessment Hypertension was described as BP  $\ge$  140 and/or  $\ge$  90 mmHg or being on drug therapy (WHO 2011 guidelines).

Stage 3: Anthropometric data collection includes Body mass index (BMI), Waist-hip ratio Height, and body mass was determined using a stadiometer and scales respectively.

#### **Statistics**

The data obtained from the Feedback form analysis was entered into Microsoft Excel 2013. Microsoft Excel and Statistical Package for Social Science (SPSS) software program (IBM SPSS v.20 Inc., Chicago II, USA) were used for descriptive analysis of the data. The results were presented as percentages and frequency distribution. The Chisquare test was utilized to evaluate the relationship between various categorical variables. P-value< 0.05 was regarded considerable.

## **RESULTS AND DISCUSSION**

**Table 1**: show the sociodemographic characteristics of participants, the study included 213 participants more than half of them 57.7% 20-30 years old, 63.8% females, and 36.2% males. 62.9% were single and 34.7% were married. 52.1% were healthy weight, 29.1% overweight, and 9.4% were obese.43.7% were students and 38.5% were the employee. As regards educational level 81.2% had collegiate. 39.9% of participants had a monthly income of more than 2000.

Table 2: show the prevalence of CVS risk factors and physical activity, 72.3% of participants joined the club for general health and fitness, 33.8% to lose weight, and 22.1% to spend spare time. The majority (95.8%) of participants have not been diagnosed with heart disease, a third of them (31.5%) are worried about getting heart disease but 68.5% know that heart disease can be prevented and 26.8% have a family history of heart disease. Only 16% reported a family member who died of heart disease in 32.4% of them the family member was grand and 61.8% reported that the person who died of heart disease was over 60 years. Only 15% were smokers of them 37.5% had been smoking for 6-10 years. 67.1% reported that they eat healthy meals but 54.5% prefer fast food and 43.7% said that they eat fruits and vegetables daily. 78.4% had an idea about an unhealthy diet, 81.7% had information about obesity and overweight, 69.5% tried to lose weight and only 18.8% suffer from a chronic disease. 28.2% of participants reported low-intensity physical activity most weeks and 22.1% had high-intensity physical activity three or more times a week. Only 10.8% had been diagnosed with high blood pressure, 56.5% from them taking prescription medications to lower blood pressure, 30.4% follow a special

diet to lower their blood pressure and 82.6% monitor their own blood pressure. 13.6% had been diagnosed with an increased level of body fat, and 37.9% of them follow a special diet to lower blood cholesterol. 8.5% of participants had been diagnosed with diabetes, 33.3% of them administrate insulin as a treatment, and the same percentage for oral antidiabetic medications. 43.7% of participants had a family history of diabetes and for women, only 12.5% had a history of gestational diabetes. 40.8% of participants reported they were under pressure all the time.

**Table 3**: show the knowledge of participants about CVD risk factors and their prevention, most participants 89.2% correctly answered that practicing physical activity can reduce the risk of cardiovascular disease, and nearly half of participants 55.4% know that eating a lot of red meat increases the risk of cardiovascular disease, 54% said that high blood pressure is defined as 120/80 or higher (systolic/diastolic).

As shown in **Table 4**, 70.9% know that eating a diet rich in fibers lowers blood cholesterol also 69.5% said that Eating fruits and vegetables prevents cardiovascular diseases. 84% said that walking can prevent the risk of cardiovascular disease and 68.5% said that usual physical activity for 150 minutes a week will reduce your chance of having a heart attack or stroke. 76.5% of participants said that being overweight or obese makes them more likely to develop cardiovascular disease, 56.3% know that people with diabetes are more at risk of having a heart attack or stroke and nearly third 36.2% said that trans fats are healthier for the heart than most other types of fats.

Table 1.	Sociodemographic (n=213)	characteristic	s of		
Parameter No. %					
	Less than 20 years	27	12.7		
	20 - 30	123	57.7		
Age	31 - 40	51	23.9		
	41 -50	8	3.8		
	51-60	4	1.9		
Condon	Male	77	36.2		
Gender	Female	136	63.8		
AT .1 11.	Saudi	211	99.1		
Nationality	Non-Saudi	2	.9		
	under weight	20	9.4		
DMI	healthy weight	111	52.1		
DIVII	over weight	62	29.1		
	obese	20	9.4		
	Student	93	43.7		
	female teacher	1	.5		
Occupation	employee	82	38.5		
	free work	10	4.7		
	I do not work	27	12.7		
	primary	2	.9		
Educational leve	el secondary	20	9.4		
	diploma	3	1.4		
	collegiate	173	81.2		

	high degree	15	7.0
	2000>	85	39.9
Monthly income	2000-7000	43	20.2
	7000<	85	39.9
Marital status	Single	134	62.9
	married	74	34.7
	Divorced	5	2.3

Table	2.	Determinants	of	cardiovascular	disease
among	pai	rticipants (n=21	3)		

Parameter			%
	to lose weight	72	33.8
Reason to join the gym	General health and fitness	154	72.3
	spending spare time	47	22.1
Diagnosed with heart disease	yes	9	4.2
Diagnosed with heart disease	no	204	95.8
Womind about active beaut disease	yes	67	31.5
worried about getting heart disease	no	146	68.5
Heart disease can be prevented	yes	146	68.5
Heart disease can be prevented	no	67	31.5
Family history of heart disease	yes	57	26.8
r anny mistory of heart disease	no	156	73.2
Family member who died of heart	yes	34	16.0
disease	no	179	84.0
	the father	10	29.4
	Grand	11	32.4
If the answer is "yes", what is the	Brother	1	2.9
relationship	Son	1	2.9
	maternal uncle	7	20.6
	paternal uncle	4	11.8
	less than 20	1	2.9
	30-39	6	17.6
Age of the person who died of	40 - 49	5	14.7
neart disease	50 - 59	1	2.9
	over 60	21	61.8
	Yes smoker	32	15.0
Smoking status	non smoker	172	80.8
	Ex smoker	9	4.2
	15th	9	28.1
	6 - 10	12	37.5
If the answer is "yes", how long	11-15	8	25.0
	16 - 20	2	6.3
	more than 20	1	3.1

# **Table 3.** Special health problems among participants(n= 213)

	/			
	Parame	ter	No.	%
	Eat haalthy maala	yes	143	67.1
Eat healthy meals		no	70	32.9
	Prefer fast food	yes	116	54.5

Eat fruits and vegetables dailyyes9343.7no12056.3Have an idea about an unhealthy dietyes16778.4Noyes17481.7obesity and overweightno3918.3Tried to lose weightyes14869.5Suffer from a chronic diseaseyes4018.8no17381.210.018.3Suffer from a chronic diseaseyes4018.8no17381.210.112.2Generally activeHigh-intensity physical activity most weeks32.212.1Generally activeHigh-intensity physical activity most weeks34.110.0Diagnosed with high blood pressureyes23.114.1Ji ft he answer is "yes" taking following a special diet to lower blood pressureyes13.114.1Ji the answer is "yes" taking hyperlipidaemiano14.115.515.5If the answer is "yes" taking lower blood pressureno16.015.616.0If the answer is "yes" taking lower blood pressureno16.615.616.0If the answer is "yes" taking long16.016.016.016.0If the answer is "yes" taking long16.016.016.016.0If the answer is "yes", follow long116.015.016.0If the answer is "yes", follow long116.016.016.0If the answer is "yes", followa		no	97	45.5
no12056.3Have an idea about an unhealthy dietyes16778.4no16778.4Have information about obesity and overweightno3918.3Tried to lose weightyes14869.5Suffer from a chronic diseaseyes4018.8no17381.2Suffer from a chronic diseaseyes4018.8no17381.2Generally activeHigh-intensity physical activity once or twice a week4219.7Generally activeHigh-intensity physical activity three or more times a week3210.8Jagnosed with high blood pressureyes2310.8If the answer is "yes" taking prescription medications to lower blood pressureyes1314.1Jiganosed with hysical activityyes1356.556.5If the answer is "yes" taking prescription medications to lower blood pressureno1669.6If the answer is "yes" taking prescription medications to lower blood pressureno1669.6If the answer is "yes", for how long1167.529If the answer is "yes", for how long291.47.41.41.47.4Jiganosed with hyperlipidaemiayes131.67.52.591.67.61.67.61.67.61.67.61.67.61.67.61.67.61.6 <t< td=""><td>Eat fruits and vegetables daily</td><td>yes</td><td>93</td><td>43.7</td></t<>	Eat fruits and vegetables daily	yes	93	43.7
Have an idea about an unhealthy dietyes16778.4no4621.6Have information about obesity and overweightno3918.3Tried to lose weightno6530.5Suffer from a chronic diseaseyes4018.8no17381.281.2Suffer from a chronic diseaseyes4018.8no17381.210.1Generally activeHigh-intensity physical activity once or twice a weeks4219.7Generally activeLi don't know/not sure3416.0There is no weekly physical activity thee or more times a week3014.1Diagnosed with high blood pressureyes2310.8If the answer is "yes" taking prescription medications to lower blood pressureyes1356.5If the answer is "yes" taking prescription medications to lower blood pressureyes1356.5If the answer is "yes" taking prescription medications to lower blood pressureyes1414.14If the answer is "yes", for how long116.67.8If the answer is "yes", for how long1167.514.15If the answer is "yes", for how long1167.514.15If the answer is "yes", for how long11167.5If the answer is "yes", for how long11167.5If the answer is "yes", for how long11167.5 </td <td></td> <td>no</td> <td>120</td> <td>56.3</td>		no	120	56.3
no4621.6Have information about obesity and overweightyes17481.7Tried to lose weightno6530.5Suffer from a chronic diseaseyes14869.5no17381.210.0173Suffer from a chronic diseaseyes6028.2activity most weeks6028.221.1Generally activeHigh-intensity physical activity once or twice a week4722.1Generally activeHigh-intensity physical activity once or twice a week4722.1Diagnosed with high blood pressureyes2310.81yes2310.810.910yes2310.82311942.222315yes1022916the answer is "yes" taking prescription medications to lower blood pressureno1043.516the answer is "yes" taking prescription medications to lower blood pressureno1669.617biagnosed with hyperlipidaemiayes1982.616the answer is "yes", for how long32916the answer is "yes", for how long32916the answer is "yes", for how long1167.516the answer is "yes", for how long1167.516the answer is "yes", for how long1167.516th	Have an idea about an unhealthy diet	yes	167	78.4
Have information about obesity and overweightyes17481.7Tried to lose weightno3918.3Tried to lose weightyes14869.5Suffer from a chronic diseaseyes4018.8no17381.2no173Only low intensity physical activity most weeks6028.2Generally activeHigh-intensity physical activity once or twice a week4219.7Generally activeHigh-intensity physical 		no	46	21.6
obesity and overweightno3918.3Tried to lose weightyes14860.5Suffer from a chronic diseaseyes4018.8no17381.2activity most weeks6028.2Generally activeHigh-intensity physical activity three or more times a week4221.1Generally activeLidon't know/not sure3416.0There is no weekly physical activity three or more times a week3014.1Diagnosed with high blood pressureyes2310.8If the answer is "yes" taking robow blood pressureyes1356.5If the answer is "yes" taking bloom pressureyes1356.5If the answer is "yes" taking hyperipidaemiayes1356.5If the answer is "yes" taking blood pressureyes1982.6If the answer is "yes" taking blood pressureyes18.45.5If the answer is "yes" taking blood pressureyes1982.6If the answer is "yes" taking blood pressureno1665.6If the answer is "yes", for how long31.47.4If the answer is "yes", for how long1167.5If the answer is "yes", for how long291.4If the answer is "yes", for how long1167.5If the answer is "yes", for how long291.4If the answer is "yes", for how long1116I	Have information about	yes	174	81.7
Tried to lose weightyes14860.5Suffer from a chronic diseaseyes4018.8no17381.2no17381.2only low intensity physical activity most weeks6028.2Generally activeHigh-intensity physical activity once or twice a week7021.1Generally activeLi don't know/not sure3416.0There is no weekly physical activity once or twice a week3614.1Diagnosed with high blood pressureyes2310.81yes2310.810.1194.22.110.1194.22.110.111010.110.110.111010.110.110.111010.110.110.111010.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.110.110.110.1110.11	obesity and overweight	no	39	18.3
Suffer from a chronic diseaseno6530.5Suffer from a chronic diseaseyes4018.8no17381.2Only low intensity physical activity once or twice a weeks6028.2High-intensity physical activity once or twice a weeks4219.7Generally activeHigh-intensity physical activity once or twice a weeks4721.1Diagnosed with high blood pressureyes2310.8Diagnosed with high blood pressureyes2310.81yes231.452194.2194.221yes, how long31.4541.910216 the answer is "yes" taking pressureyes1356.517 the answer is "yes" taking pressureyes1669.616 the answer is "yes" taking pressureyes1982.617 the answer is "yes" taking pressureyes1982.616 the answer is "yes", for how long1167.517 the answer is "yes", for how long2931.452931.41.616 the answer is "yes", follow a special diet to lower blood cholesterol1167.517 the answer is "yes", follow a special diet to lower blood cholesterolyes188.518 the answer is "yes", follow a special diet to lower blood cholesterolyes18<	Tried to lose weight	yes	148	69.5
Suffer from a chronic diseaseyes4018.8no17381.2no17381.2Only low intensity physical activity most weeks6028.2High-intensity physical activity once or twice a week4219.7Generally activeHigh-intensity physical activity three or more times a week472.1I don't know/not sure3416.0There is no weekly physical activity3014.1Diagnosed with high blood pressureyes2310.81942.252.3194.2252194.1194.12194.12194.231541.9102.931165.541165.511669.61167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.51167.5116<		no	65	30.5
no17381.2Only low intensity physical activity most weeks6028.2High-intensity physical activity once or twice a week4219.7Generally activeHigh-intensity physical activity three or more times a week4722.1Diagnosed with high blood pressureyes2310.8Diagnosed with high blood pressureyes2310.8194.225194.225194.225194.225191.454102.91.4102.91.41194.2.91541.91021610yes1.356.517991.43.51610yes1.91043.517102.91.43.616101043.5.5.51710101.6.5.6181116.7.5.6.2191116.7.5.911101.4.5.2.9141116.7.5.2.9161116.7.5.2.9161116.7.5.2.9161116.7.5.2.	Suffer from a chronic disease	yes	40	18.8
Only low intensity physical activity most weeks6028.2High-intensity physical activity once or twice a week4219.7Generally activeHigh-intensity physical activity three or more times a week4721.1Diagnosed with high blood pressureyes2310.8Diagnosed with high blood pressureyes2310.8194.222194.222194.22311029102194.11029194.2252.31101029102110104.110104.1110104.1104.1110104.1104.1110104.11.11.6110101.11.61.11101.11.61.11.61101.21.11.61.21101.21.11.61.21111.61.21.411.61.41.41.411.61.41.411.61.41.411.61.41.411.61.41.411.61.41.41 <td></td> <td>no</td> <td>173</td> <td>81.2</td>		no	173	81.2
High-intensity physical activity once or twice a week429.7Generally activeHigh-intensity physical activity three or more times a week472.1I don't know/not sure3014.0There is no weekly physical activity3014.1Diagnosed with high blood pressureyes2310.8If yes, how long33.114.1If yes, how long33.114.1If the answer is "yes" taking prescription medications to lower blood pressureyes1356.5If the answer is "yes" monitor own blood pressureyes102330.1If the answer is "yes" monitor own blood pressureyes1982.6If the answer is "yes" monitor own blood pressureyes1982.6If the answer is "yes", for how long1167.5If the answer is "yes", for how long1167.5If the answer is "yes", for how long1167.5If the answer is "yes", for how long1111If the answer is "yes", for how long1131.4If the answer is "yes", for how long1131.4If the answer is "yes", for how long1131.4If the answer is "yes", for how long111.51.5If the answer is "yes", for how long111.51.5If the answer is "yes", for how long11 <td< td=""><td></td><td>Only low intensity physical activity most weeks</td><td>60</td><td>28.2</td></td<>		Only low intensity physical activity most weeks	60	28.2
Generally activeHigh-intensity physical activity three or more times a week4722.1I don't know/not sure3416.0There is no weekly physical activity3014.1Diagnosed with high blood pressureyes2310.8194.2252.3If yes, how long331.4541.9102.9If the answer is "yes" taking prescription medications to lower blood pressureyes1356.51f the answer is "yes" taking volod pressureyes1356.51f the answer is "yes" monitor own blood pressureyes1982.61f the answer is "yes" monitor own blood pressureyes1982.61f the answer is "yes" monitor own blood pressureyes1982.61f the answer is "yes", for how long1167.51f the answer is "yes", for how long1167.51f the answer is "yes", for how long1167.51f the answer is "yes", follow a special diet to lower blood cholesterolyes1137.91f the answer is "yes", follow a special diet to lower blood cholesterolyes188.51101315.915.51101315.915.511013.515.915.511167.515.511167.515.511		High-intensity physical activity once or twice a week	42	19.7
I don't know/not sure3416.0There is no weekly physical activity3014.1Diagnosed with high blood pressureyes2310.8no1089.2194.2252.31.4194.2194.2252.3If yes, how long331.41029102910291110101043.516.51110101043.516.511167.5936.516.511169.6101043.511169.61016.516.511169.610101011169.61016.516.511169.61016.516.511169.616.516.516.511167.529.516.511167.529.516.511167.529.516.511167.529.516.511167.529.516.511167.529.516.511167.529.516.511167.516.516.516.511167.516.516.	Generally active	High-intensity physical activity three or more times a week	47	22.1
There is no weekly physical activity3014.1Diagnosed with high blood pressureyes2310.8no1089.21252.31252.31541.9251.01.0541.91.010291194.2541.9102910291110311104.311101.011101.011101.011101.0111.01.0121.01.0 <t< td=""><td></td><td>I don't know/not sure</td><td>34</td><td>16.0</td></t<>		I don't know/not sure	34	16.0
$\begin{array}{llllllllllllllllllllllllllllllllllll$		There is no weekly physical activity	30	14.1
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Diagnosed with high blood	yes	23	10.8
If yes, how long         1         9         4.2           If yes, how long         3         3         1.4           5         4         1.9           10         2         .9           If the answer is "yes" taking prescription medications to lower blood pressure         no         10         43.5           If the answer is "yes" taking prescription medications to lower blood pressure         no         10         43.5           If the answer is "yes" monitor lower blood pressure         no         16         69.6           If the answer is "yes" monitor own blood pressure         no         4         17.4           Diagnosed with hyperlipidaemia         yes         29         13.6           If the answer is "yes", for how long         3         2         .9           If the answer is "yes", for how long         3         2         .9           If the answer is "yes", for how long         3         2         .9           If the answer is "yes", for how long         3         2         .9           If the answer is "yes", for how long         3         1         .4           5         2         .9         .9         .1           If the answer is "yes", for how long         yes         .1	pressure	no	190	89.2
If yes, how long         3         3         1.4           5         4         1.9           10         2         .9           If the answer is "yes" taking prescription medications to lower blood pressure         no         10         43.5           If the answer is "yes" taking prescription medications to lower blood pressure         no         10         43.5           If the answer is "yes" monitor lower blood pressure         no         10         43.5           If the answer is "yes" monitor own blood pressure         no         16         69.6           If the answer is "yes" monitor own blood pressure         no         18         82.6           Diagnosed with hyperlipidaemia         no         184         86.4           Diagnosed with hyperlipidaemia         no         184         86.4           If the answer is "yes", for how long         3         2         .9           If the answer is "yes", for how long         3         1.4         .5         .2           If the answer is "yes", for how long         yes         .1         .6         .2.8           If the answer is "yes", for how long         yes         .1         .3.7         .2         .9         .3         .2         .9         .9         .3 <t< td=""><td></td><td>1</td><td>9</td><td>4.2</td></t<>		1	9	4.2
If yes, how long331.4541.9102.9If the answer is "yes" taking prescription medications to lower blood pressureno1043.5If the answer is "yes" rescription medications to lower blood pressureno1043.5If the answer is "yes" monitor own blood pressureyes730.4If the answer is "yes" monitor own blood pressureyes1982.6If the answer is "yes" monitor own blood pressureyes2913.6Diagnosed with hyperlipidaemiayes2913.6If the answer is "yes", for how long1167.5262.81.452If the answer is "yes", for how long32.9431.452.9If the answer is "yes", follow a special diet to lower blood cholesterolyes1137.9If the answer is "yes", follow a special diet to lower blood cholesterolyes188.5Diagnosed with diabetesyes188.5.91.5If the answer is "yes", what is treatment for diabetesYes188.5.91.5If the answer is "yes", what is treatment for diabetes.61 antidiabetic medications6.33.3If the answer is "yes", what is treatment for diabetes.61 antidiabetic medications6.33.3If the answer is "yes", what is treatment for diabetes.61 antidiabetic medications6.33.3 </td <td></td> <td>2</td> <td>5</td> <td>2.3</td>		2	5	2.3
	If yes, how long	3	3	1.4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		5	4	1.9
If the answer is "yes" taking prescription medications to lower blood pressureyes1356.5If the answer is "yes"no1043.5If the answer is "yes"yes730.4following a special diet to lower blood pressureno1669.6If the answer is "yes" monitor own blood pressureyes1982.6If the answer is "yes" monitor own blood pressureyes2913.6Diagnosed with hyperlipidaemiayes2913.6If the answer is "yes", for how long1167.5262.82.99431.45252.9431.452.9137.9special diet to lower blood cholesterolno1862.1Diagnosed with diabetesyes188.5no19591.5.5.5If the answer is "yes", what is treatment for diabetesOral antidiabetic medications Self-monitoring of blood sugar211.1following a diet422.2.5.5		10	2	.9
prescription medications to lower blood pressure no 10 43.5 If the answer is "yes" yes 7 30.4 following a special diet to lower blood pressure no 16 69.6 If the answer is "yes" monitor own blood pressure no 4 17.4 Diagnosed with hyperlipidaemia no 184 86.4 I 1 16 7.5 If the answer is "yes", for how long 3 2 9 4 3 1.4 5 2 9 11 37.9 If the answer is "yes", follow a special diet to lower blood cholesterol no 18 82.6 no 184 86.4 I 1 16 7.5 2 9 4 3 1.4 5 2 9 4 3 1.4 5 9 11 37.9 If the answer is "yes", follow a special diet to lower blood cholesterol no 18 82.1 Diagnosed with diabetes yes 18 8.5 no 195 91.5 insulin 6 33.3 If the answer is "yes", what is treatment for diabetes 2 11.1 following a diet 4 22.2	If the answer is "yes" taking	yes	13	56.5
Index <th< td=""><td>prescription medications to</td><td>20</td><td>10</td><td>13 5</td></th<>	prescription medications to	20	10	13 5
If the answer is "yes" is yes is a special diet to lower blood pressure is "yes" is non it is a special diet to lower blood pressure is "yes" monitor is no is a special diet to own blood pressure is "yes" is "yes		lio	10	45.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	following a special diet to	yes	1	50.4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	lower blood pressure	no	16	69.6
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	If the answer is "yes" monitor	yes	19	82.6
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	own blood pressure	no	4	17.4
hyperlipidaemia no 184 86.4 hyperlipidaemia no 184 86.4 1 16 7.5 2 6 2.8 1 16 7.5 2 9 4 3 1.4 5 2 9 4 3 1.4 5 2 9 1 37.9 If the answer is "yes", follow a yes 11 37.9 picture to lower blood cholesterol no 18 62.1 Diagnosed with diabetes yes 18 8.5 no 195 91.5 insulin 6 33.3 If the answer is "yes", what is treatment for diabetes Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2	Diagnosed with	yes	29	13.6
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	hyperlipidaemia	no	184	86.4
If the answer is "yes", for how long $\begin{array}{cccccccccccccccccccccccccccccccccccc$		1	16	7.5
long 3 2 .9 4 3 1.4 5 2 .9 If the answer is "yes", follow a special diet to lower blood cholesterol no 18 62.1 Diagnosed with diabetes yes no 18 8.5 no 195 91.5 insulin 6 33.3 If the answer is "yes", what is treatment for diabetes 6 33.3 Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2	If the answer is "yes", for how	2	6	2.8
431.452.9If the answer is "yes", follow a special diet to lower blood cholesterolyes1137.9Diagnosed with diabetesno1862.1Diagnosed with diabetesyes no188.5 19591.5If the answer is "yes", what is treatment for diabetesOral antidiabetic medications Self-monitoring of blood sugar633.3If oflowing a diet422.22	long	3	2	.9
If the answer is "yes", follow a special diet to lower blood cholesterol no 18 62.1 Diagnosed with diabetes yes 18 8.5 no 195 91.5 insulin 6 33.3 If the answer is "yes", what is treatment for diabetes Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2		4	3	1.4
If the answer is "yes", follow a no 18 62.1 Diagnosed with diabetes yes no 19 91.5 If the answer is "yes", what is treatment for diabetes Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2	TC.1	ves	2 11	.9 37.9
Diagnosed with diabetesyes no18 1958.5 91.5If the answer is "yes", what is treatment for diabetesOral antidiabetic medications Self-monitoring of blood sugar633.3If the answer is "yes", what is treatment for diabetesOral antidiabetic medications Self-monitoring of blood sugar632.3If the answer is "yes", what is treatment for diabetesOral antidiabetic medications Self-monitoring of blood sugar623.3If the answer is treatment for diabetesIf the answer is treatment for diabetes211.1If the answer is treatment for diabetes422.2	If the answer is "yes", follow a special diet to lower blood cholesterol	no	18	62.1
insulin 6 33.3 If the answer is "yes", what is treatment for diabetes Cral antidiabetic medications 6 33.3 Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2	Diagnosed with diabetes	yes no	18 195	8.5 91.5
If the answer is "yes", what is treatment for diabetes Oral antidiabetic medications 6 33.3 Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2		insulin	6	33.3
Self-monitoring of blood sugar 2 11.1 following a diet 4 22.2	If the answer is "yes", what is	Oral antidiabetic medications	6	33.3
following a diet 4 22.2	treatment for diabetes	Self-monitoring of blood sugar	2	11.1
		following a diet	4	22.2

Family history of diabetes	yes	93	43.7
	no	120	56.3
History of gestational diabetes	yes	17	12.5
in women	No	119	87.5
Under pressure all the time	yes	87	40.8
ender pressure un the time	No	126	59.2

## Table 4. Knowledge of participants of risks and prevention of cardiovascular disease (n=213)

Parameter	Yes	No	Don't know
You can reduce the risk of cardiovascular disease by practicing physical activity	190	5	18
	89.2%	2.3%	8.5%
Eating a lot of red meat increases the risk of cardiovascular disease	118	31	64
	55.4%	14.6%	30.0%
High blood pressure is defined as 120/80 or	115	42	56
higher (systolic/diastolic)	54.0%	19.7%	26.3%
Eating a diet rich in fibre lowers blood cholesterol	151	13	49
	70.9%	6.1%	23.0%
Eating fruits and vegetables prevents cardiovascular diseases	148	19	46
	69.5%	8.9%	21.6%
Walking can prevent the risk of cardiovascular disease	179	7	27
	84.0%	3.3%	12.7%
Usual physical activity for 150 minutes a week will reduce your chance of having a heart attack or stroke	146 68.5%	16 7.5%	51 23.9%
People with diabetes are more at risk of having	120	23	70
a heart attack or stroke	56.3%	10.8%	32.9%
Trans fats are healthier for the heart than most other types of fats	77	56	80
	36.2%	26.3%	37.6%
You are more likely to develop cardiovascular disease if you are overweight or obese.	163	15	35
	76.5%	7.0%	16.4%

Cardiovascular diseases (CVDs) are a group of illnesses associated with heart and blood vessels. (CVDs) are common in the general population, mainly among adults [17]. Based on the WHO, heart disease more specifically coronary heart disease is the major cause of death worldwide and among the major health burdens globally and more than 17 million people die annually (30% of all deaths) [17]. According to various epidemiological studies, many nonmodifiable and modifiable CVDs risk factors were recognized. Old age, family history, and male gender are nonmodifiable risk factors, but modifiable risk factors comprises smoking, hypertension, diabetes, dyslipidemia, obesity, physical inactivity, unhealthy diets, and heavy alcohol intake [18]. One study indicates the prevalence of CVD mortality rate in the Kingdom of Saudi Arabia is 46% [19]. This significantly high percentage might be related to the lack of awareness of cardiovascular risk factors. Encourage people to adopt a cardio-protective lifestyle. Little is known about cardiovascular risk factors in the general population. Knowledge is an essential step in developing a cardiovascular protective lifestyle [20]. This is a cross-sectional prospective study conducted among 213 unselected adult populations attending the Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia. The study aims at determining the incidence and awareness of cardiovascular disease risk factors among the adult population attending the Fitness center/Gymnasium in Arar, Kingdom of Saudi Arabia.

According to national studies [21-23] the incidence rates of CVDs risk factors are disturbing drift over time. Smoking is said to cause almost ten percent of all CVDs then physical inactivity (6%), as well as overweight and obesity (5%) [24]. An increase in BP levels also increases the risk of stroke and coronary heart disease [25]. Moreover, individuals with diabetes have a two-fold risk of vascular disease, and surged cholesterol and triglyceride levels are independent risk factors for coronary heart disease [26, 27]. Individual knowledge of cardiovascular disease risk factors is paramount in combating these factors. However, few studies worldwide have addressed knowledge of cardiovascular disease risk factors. As regards awareness of cardiovascular disease risk factors and their prevention among participants, results from our study showed that nearly half of participants 55.4% know that eating a lot of red meat increases the risk of cardiovascular disease, 54% said that high blood pressure is defined as 120/80 or higher (systolic/diastolic), 70.9% know that eating a diet rich in fibers lowers blood cholesterol also 69.5% said that Eating fruits and vegetables prevents cardiovascular diseases. 76.5% of participants said that being overweight or obese makes them more likely to develop cardiovascular disease, 56.3% know that people with diabetes are more at risk of having a heart attack or stroke and nearly third 36.2% said that trans fats are healthier for the heart than most other types of fats. In Egypt, another study conducted among 200 participants reported that 59% of them know that eating a lot of red meat increases heart disease risk, 41% reported stress as the most important cause of heart attacks, 44% think that high blood cholesterol is a risk factor for heart diseases, 37% think that eating a high fibers diet increases the risk of getting heart disease, 28% said that obese people are more at risk for getting heart disease and only 13% said that elevation of blood sugar for a long time is a risk factor for heart disease [28]. In India, another study carried out among the adult population found that all of the participants said that being overweight increases a person's risk for heart disease followed by 98% knowing that high cholesterol is a risk factor for developing heart disease, 94% reported High Blood pressure as a risk factor for heart disease and 92% reported Smoking as a risk factor for heart disease [29]. In Nepal, a descriptive cross-sectional study included 200 adults reported that most of the respondents (90.5%) were aware of smoking as a risk factor for CVD, 82.5% know that high blood pressure is a risk factor for heart disease, 81% said that daily intake of fruits and vegetables reduces the risk for developing heart disease, 79.5% think that stress and anxiety increase the risk of heart disease, 78.5% said that keeping blood pressure under control will reduce a person's risk for developing heart disease, 72.5% of participants reported that being overweight

increases a person's risk for heart disease, 60.5 % said that high cholesterol is a risk factor for developing heart disease and 59.5% know that diabetes is a risk factor for developing heart disease [30]. In Cameron, another study reported that participants were aware that smoking (82%), unhealthy diet (70.6%), lack of exercise (67.0%), obesity (69.7%), stress (73.1%), high blood pressure (HBP) (73.3%) and diabetes (60.8%) were potential risk factors for CVD [31]. Physical activity (PA) and exercise training play a central role in maintaining good health and preventing primary and secondary diseases. Gyms and fitness centers are currently largely reachable in large city and several smaller towns and have a critical role in employing strategies to decrease the population's CV risk factors [32]. Multiple physiological mechanisms are involved, and the impact of PA on reducing cardiovascular (CV) risk factors is well established across different exercise modalities [33]. Recent first-line management of arterial hypertension, hypercholesterolemia, obesity, diabetes, and non-alcoholic fatty liver disease depends on dietary management, lifestyle modification, and, particularly, physical exercise [34, 35]. In accordance with this, our study reported that most participants 89.2% correctly answered that practicing physical activity can reduce the risk of cardiovascular disease, 84% said that walking can prevent the risk of cardiovascular disease and 68.5% said that usual physical activity for 150 minutes a week will reduce your chance of having a heart attack or stroke. Similar to our results another study reported that 42% of participants know that only exercise in a GYM decrease risk of heart disease and 34% said that regular exercises decrease the risk of heart disease [28]. Also, in India, results from another study show that 92% of participants correctly answered that only exercising at a gym or in a class will help lower a person's chance of developing heart disease and 90% said that regular Physical activity will person's chance of getting heart disease [29]. In Nepal, another study found that 86.5% know that regular physical activity will lower a person's chance of getting heart disease, and 82.5% said that walking and gardening are considered an exercise that will help lower a person's chance of developing heart disease, however, half of the participants (50%) were aware of only exercising at a gym will not help lower person's chance of developing heart disease [30].

According to the prevalence of CVD risk factors among participants, our study reported that the majority (95.8%) of participants have not been diagnosed with heart disease but 26.8% have a family history of heart disease, only 15% were smokers, 67.1% reported that they eat healthy meals and 43.7% said that they eat fruits and vegetables daily. 28.2% of participants reported low-intensity physical activity most weeks and 22.1% had high-intensity physical activity three or more times a week. Only 10.8% had been diagnosed with high blood pressure, 13.6% had been diagnosed with an increased level of body fat and 8.5% had been diagnosed with diabetes. In Nepal, another study reported that most of the participants (89.5%) of them had unhealthy dietary practices, only 10.5% had adequate consumption of fruits and

vegetables daily, 11.5% were current smokers, about physical activity, almost (96.5%) of respondents were physically active which contrasted with our results, 60.5% had normal BMI, only 12.5% were hypertensive and 11.5% had a family history of CVD [30]. In Lebanon, another study found that 42% of the studied population reported smoking either cigarettes or waterpipe only 2.1% reported heavy alcohol consumption of ≥3 drinks/day, and 17.3% and 8.7% reported not consuming at all fruits and vegetables, respectively. However, the prevalence of moderate exercise and walking was high, were 54.5%, and 70% reported engaging in moderate activities and walking for at least 10 minutes for  $\geq 4$ times/week, 46.5% of the study participants were overweight and 29.4% were obese and the prevalence of reported hypertension, diabetes, and dyslipidemia previously diagnosed by a doctor was 29.8% 22.8% and 22.5% respectively [36].

## CONCLUSION

The present study concluded that there was good awareness of cardiovascular disease risk factors among participants which was in the same line with other reported studies. Also, they know the role of physical activity in reducing the risk of cardiovascular disease risk.

ACKNOWLEDGMENTS: Many thanks to Pr. Anshoo Agarwal; Professor of Pathology, Faculty of Medicine, Northern Border University, Arar, Saudi Arabia, for his continuous help, support and encouragement to complete this work.

CONFLICT OF INTEREST: None FINANCIAL SUPPORT: None

ETHICS STATEMENT: Ethical approval was obtained from the research ethics committee of Northern Border University.

## References

- 1. Guo L, Zhang S. Association between ideal cardiovascular health metrics and risk of cardiovascular events or mortality: a meta-analysis of prospective studies. Clin Cardiol. 2017;40(12):1339-46.
- Gaye B, Canonico M, Perier MC, Samieri C, Berr C, Dartigues JF, et al. Ideal Cardiovascular Health, Mortality, and Vascular Events in Elderly Subjects: The Three-City Study. J Am Coll Cardiol. 2017;69(25):3015-26.
- Enserro DM, Vasan RS, Xanthakis V. Twenty-Year Trends in the American Heart Association Cardiovascular Health Score and Impact on Subclinical and Clinical Cardiovascular Disease: The Framingham Offspring Study. J Am Heart Assoc. 2018;7(11):e008741.
- Yeboah J. Road to the American Heart Association 2020 Impact Goals: The Metric for Monitoring Progress. Circ Cardiovasc Imaging. 2018;11(1):e007385.
- Kivimäki M, Batty GD, Hamer M, Ferrie JE, Vahtera J, Virtanen M, et al. Using additional information on working hours to predict coronary heart disease: a cohort study. Ann Intern Med. 2011;154(7):457-63.
- Kivimäki M, Batty GD, Singh-Manoux A, Britton A, Brunner EJ, Shipley MJ. Validity of cardiovascular disease event ascertainment using linkage to UK hospital records. Epidemiology. 2017;28(5):735-9.
- Britton A, Milne B, Butler T, Sanchez-Galvez A, Shipley M, Rudd A, et al. Validating self-reported strokes in a longitudinal UK cohort study (Whitehall II): Extracting information from hospital medical records versus the Hospital Episode Statistics database. BMC Med Res Methodol. 2012;12:83.

- Gaye B, Prugger C, Perier MC, Thomas F, Plichart M, Guibout C, et al. High level of depressive symptoms as a barrier to reach an ideal cardiovascular health. The Paris Prospective Study III. Sci Rep. 2016;6:18951.
- Kronish IM, Carson AP, Davidson KW, Muntner P, Safford MM. Depressive symptoms and cardiovascular health by the American Heart Association's definition in the Reasons for Geographic and Racial Differences in Stroke (REGARDS) study. PLoS One. 2012;7(12):e52771.
- Huffman MD, Capewell S, Ning H, Shay CM, Ford ES, Lloyd-Jones DM. Cardiovascular health behavior and health factor changes (1988-2008) and projections to 2020: results from the National Health and Nutrition Examination Surveys. Circulation. 2012;125(21):2595-602.
- Shah AM, Claggett B, Folsom AR, Lutsey PL, Ballantyne CM, Heiss G, et al. Ideal Cardiovascular Health During Adult Life and Cardiovascular Structure and Function Among the Elderly. Circulation. 2015;132(21):1979-89.
- Aatola H, Hutri-Kähönen N, Juonala M, Laitinen TT, Pahkala K, Mikkilä V, et al. Prospective relationship of change in ideal cardiovascular health status and arterial stiffness: the Cardiovascular Risk in Young Finns Study. J Am Heart Assoc. 2014;3(2):e000532.
- Hwang SJ, Onuma O, Massaro JM, Zhang X, Fu YP, Hoffmann U, et al. Maintenance of Ideal Cardiovascular Health and Coronary Artery Calcium Progression in Low-Risk Men and Women in the Framingham Heart Study. Circ Cardiovasc Imaging. 2018;11(1):e006209.
- Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, et al. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. Circulation. 2013;127(1):e6-245.
- 15. Barth J, Schneider S, von Kanel R. Lack of social support in the etiology and the prognosis of coronary heart disease: A systematic review and meta-analysis. Psychosom Med. 2010;72(3):229-38.
- Dale J, Williams S, Bowyer V. What is the effect of peer support on diabetes outcomes in adults? A systematic review. Diabet Med. 2012;29(11):1361-77.
- 17. Mendis S, Puska P, Norrving B, World Health Organization. Global atlas on cardiovascular disease prevention and control. World Health Organization; 2011. Available from: http://www.who.int/cardiovascular\_diseases/publications/atlas\_cvd/en /
- Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart disease and stroke statistics--2014 update: a report from the American Heart Association. Circulation. 2014;129(3):e28e292. doi:10.1161/01.cir.0000441139.02102.80
- WHO. Noncommunicable Diseases (NCD) Country profiles, 2018. WHO. Geneva. 2018. Available from: https://www.who.int/nmh/countries/sau\_en.pdf
- Potvin L, Richard L, Edwards AC. Knowledge of cardiovascular disease risk factors among the Canadian population: relationships with indicators of socioeconomic status. CMAJ. 2000;162(9 Suppl):S5-11.
- Nasreddine L, Naja FA, Sibai AM, Helou K, Adra N, Hwalla N. Trends in nutritional intakes and nutrition-related cardiovascular disease risk factors in Lebanon: the need for immediate action. J Med Liban. 2014;62(2):83-91.
- Nasreddine L, Naja F, Chamieh MC, Adra N, Sibai AM, Hwalla N. Trends in overweight and obesity in Lebanon: evidence from two national cross-sectional surveys (1997 and 2009). BMC Public Health. 2012;12(1):120-9.
- Naja F, Hwalla N, Itani L, Salem M, Azar ST, Zeidan MN, et al. Dietary patterns and odds of Type 2 diabetes in Beirut, Lebanon: a case-control study. Nutr Metab (Lond). 2012;9(1):111. doi:10.1186/1743-7075-9-111
- 24. World Health Organization. Global health risks: mortality and burden of disease attributable to selected major risks. World Health Organization; 2009.
- Whitworth JA; World Health Organization, International Society of Hypertension Writing Group. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. J Hypertens. 2003;21(11):1983-92.
- 26. Sarwar N, Gao P, Seshasai SR, Gobin R, Kaptoge S, Di Angelantonio E, et al. Emerging Risk Factors Collaboration. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a

collaborative meta-analysis of 102 prospective studies. Lancet. 2010;375(9733):2215-22.

- Nelson RH. Hyperlipidemia as a risk factor for cardiovascular disease. Prim Care. 2013;40(1):195-211.
- Mohamed HS. Knowledge of Coronary Heart Disease and Risk Factors Among Type 2 Diabetes Mellitus at Outpatient Department of Kasr Alainy Hospital. Egypt J Hosp Med. 2021;82(4):785-91.
- George C. A population-based study on Awareness of Cardiovascular Disease Risk Factors. Indian J Pharm Pract. 2014;7(2):23-5. doi:10.5530/ijopp.7.2.5
- Budhathoki R, Thulung B. Awareness regarding Cardiovascular Disease Risk Factors among Adults attending at a Teaching Hospital, Bharatpur. IOSR J Nurs Health Sci. 2021;10(4):1-8.
- Aminde LN, Takah N, Ngwasiri C, Noubiap JJ, Tindong M, Dzudie A, et al. Population awareness of cardiovascular disease and its risk factors in Buea, Cameroon. BMC Public Health. 2017;17(1):545. doi:10.1186/s12889-017-4477-3
- Booth FW, Roberts CK, Thyfault JP, Ruegsegger GN, Toedebusch RG. Role of Inactivity in Chronic Diseases: Evolutionary Insight and Pathophysiological Mechanisms. Physiol Rev. 2017;97(4):1351-402. doi:10.1152/physrev.00019.2016

- Sandbakk SB, Nauman J, Zisko N, Sandbakk Ø, Aspvik NP, Stensvold D, et al. Sedentary Time, Cardiorespiratory Fitness, and Cardiovascular Risk Factor Clustering in Older Adults--the Generation 100 Study. Mayo Clin Proc. 2016;91(11):1525-34. doi:10.1016/j.mayocp.2016.07.020
- Hansen D, Niebauer J, Cornelissen V, Barna O, Neunhäuserer D, Stettler C, et al. Exercise Prescription in Patients with Different Combinations of Cardiovascular Disease Risk Factors: A Consensus Statement from the EXPERT Working Group. Sports Med. 2018;48(8):1781-97. doi:10.1007/s40279-018-0930-4
- Negi CK, Babica P, Bajard L, Bienertova-Vasku J, Tarantino G. Insights into the molecular targets and emerging pharmacotherapeutic interventions for nonalcoholic fatty liver disease. Metabolism. 2022;126:154925. doi:10.1016/j.metabol.2021.154925
- 36. Fahs I, Khalife Z, Malaeb D, Iskandarani M, Salameh P. The Prevalence and Awareness of Cardiovascular Diseases Risk Factors among the Lebanese Population: A Prospective Study Comparing Urban to Rural Populations. Cardiol Res Pract. 2017;2017:3530902. doi:10.1155/2017/3530902