

Violence against Health Care Workers of Pediatric Departments in Saudi Arabia: Systematic Review

Hamed Salem S Albalwei^{1*}, Nazim Faisal Hamed Ahmed², Nasser Mufleh Saleh Albalawi³, Suad Salem Awad Albalawi⁴, Nujood Hamad Kashan Al-enazi⁴

¹Director of the Maternal and Child Health Center, Tabuk, Saudi Arabia. ²Department of Pediatrics, Maternal and Child Health Center, Tabuk, Saudi Arabia. ³Maternal and Child Health Center, Tabuk, Saudi Arabia. ⁴Department of Nursing, Maternal and Child Health Center, Tabuk, Saudi Arabia.

Abstract

Work Place Violence WPV is defined as the intentional use of physical or psychological force to injure, intimidate, or attack an individual in a work atmosphere. It is often known as any threat or physical abuse, assault, bullying, or other intimidating disrupting behavior that happens at the workplace. This is a systematic review was carried out, including PubMed, Google Scholar, and EBSCO that examining randomized controlled trials, observational, and experimental studies that study violence against healthcare practitioners in pediatric departments. The review included 6 randomized studies concerned about workplace violence against pediatric staff. Pediatric physicians and nurses are exposed to different types of violence from parents or caregivers of the attending children in their departments due to the psychological stress caused by the condition of their child. Recommendations to teach pediatric staff how to deal with violent events and how to report them must be instructed clearly.

Keywords: Violence, Healthcare workers, Workplace violence, Pediatrics departments, Pediatric staff, WPV

INTRODUCTION

Work Place Violence WPV is defined as the intentional use of physical or psychological force to injure, intimidate, or attack an individual in a work atmosphere. It is often known as any threat or physical abuse, assault, bullying, or other intimidating disrupting behavior that happens at the workplace [1]. It has been reported that 4.9–65% of health care professionals have been exposed to physical injury in their workplace and 1.2% of workplace homicide victims in the USA. Other studies reported a higher incidence of WPV [2].

HCWs are mostly vulnerable to reactive violence on the part of patients and their companions, affecting their physical and mental well-being, as patients and their families who are subject to mental stress due to hospitalization or disease may use violence against healthcare workers [3-6]. Job fatigue, customer demands, and declining patient-staff relationships have been linked with physical abuse against health care workers in the workplace [7].

It is categorized as physical and psychological violence. Physical abuse requires the use of physical force against someone or the use of objects to strike someone. It involves hitting, kicking, slapping, pulling, biting, pinching, wounding with sharp sticks, and sexual assault [8]. It may result in severe injury, dysfunction, permanent disability, or no harm at all. Non-physical aggression may involve bullying, slurs, threats, or sexual harassment; it does not cause physical

injuries but may cause psychological damage, such as stress, anxiety, low job satisfaction, and low work performance [9]. Also, WPV has many negative effects, resulting not only in physical consequences but also in relational consequences for healthcare workers at work, correlated with the decision to leave, stress out, and minimize career success for medical practitioners [10]. These effects of aggression in the workplace can lead to a loss in efficiency and also affect the quality of treatment. Additionally, the shortage of workers and the investment of defensive tactics induced by abuse in the workplace will practically raise health costs [11].

In outpatient clinics, pediatrics care services are now diminishing and pediatrics outpatient clinics were even

Address for correspondence: Hamed Salem S Albalwei, Director of the Maternal and child health center, Tabuk, Saudi Arabia.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to cite this article: Albalwei HSS, Ahmed NFH, Albalawi NMS, Albalawi SSA, Al-enazi NHK. Violence against Health Care Workers of Pediatric Departments in Saudi Arabia. Arch Pharm Pract. 2021;12(1):79-84. <https://doi.org/10.51847/RJ683KIDZS>

discontinued, resulting in a drastic rise in the number of outpatient clinics in children's hospitals, overworked medical personnel, inadequate quality behaviors, crowded conditions, and extended waiting times; these factors contribute to frustration and dissatisfaction among pediatrics patients' families which may lead to increased risk of violence [12]. Surveys were undertaken to examine the occurrence and intensity of physical violence toward health care workers committed by patients and tourists at the workplace. It is becoming an important research topic in Saudi Arabia, where cultural and ethical standards may vary dramatically from other locations Occupational abuse (WPV) committed by patients and tourists against nurses and doctors is a concern in adult emergency rooms (EDs), but largely unrecognized and unreported in pediatric EDs [13, 14].

Pediatric nurses working in hospital inpatient units have been under-studied in respect to their understanding of abuse in the workplace. Few studies were found in the pieces of literature that concentrate on occupational abuse against pediatrics nurses by patients and tourists [15].

Aim of the Study

Up to our knowledge; there's a lack of studies reporting violence against pediatric staff in Saudi hospitals, so the main objective of the study is to study the updates on violence against pediatric physicians and nurses.

MATERIALS AND METHODS

A systematic review was carried out, including PubMed, Google Scholar, and EBSCO using the following terms in

different combinations: violence against healthcare workers, workplace violence in pediatric departments, pediatric staff and workplace violence, WPV in Saudi Arabia. **Figure 1** is a flow chart that illustrates the data extraction process of the study. We included all full texts [randomized controlled trials, observational, and experimental studies which study violence against healthcare practitioners in pediatric departments. The authors extracted the data, and then the author's names, year, and region of publication, the study type, period of study, and the result were reported (**Table 1**).

Statistical Analysis

No software has been utilized to analyze the data. The data was extracted based on a specific form that contains (Author's name, publication year, country, methodology, and results). These data were reviewed by the group members to determine the initial findings and the modalities of performing the surgical procedure. Double revision of each member's outcomes was applied to ensure the validity and minimize the mistakes.

RESULTS AND DISCUSSION

The search of the mentioned databases returned a total of 66 studies that were included for title screening, 49 of them were included for abstract screening, which leads to the exclusion of 24 articles. The remaining 25 publications full-texts were reviewed. The full-text revision leads to the exclusion of 19 studies, and 6 were enrolled for final data extraction (**Table 1**). The included studies had different study designs.




Figure 1. Flow chart of the data extraction process of the study.

Gillespie, G. (2010) found that; both genders and all professional categories were at risk of witnessing verbal and physical WPV. Popular features of the suspects included patients undergoing a psychological examination and visitors suffering acute anxiety. Effects have been witnessed by staff, offenders, patients, and health employers [16].

Li, Zhe *et al.* (2017) found that; 68.6% of respondents experienced at least one WPV event in the past year. (94.9%) of offenders were family members of patients. Most of the WPV happened during the day shift (70.7%). Males were almost twice as females at risk to experience violence. As a consequence of WPV, aggression, anger, the decline in work, and work efficiency was reduced [17].

Shaw J. (2015) reported that; At least 26% of the workers shared worry about protection every week. Twenty-seven percent witnessed fear-causing conditions at least weekly. The primary causes of fear were patient or guest frustration (with the possibility of violence) and ED weapons. Respondents would be safer" with increased involvement of hospital security personnel (55%) and local police officers (71%) [18].

Strollo, Bonnie A., *et al.* (2011) in New York suggested that there is violence in the workplace and a great deal needs to be done to deter and control violence in the workplace toward nurses who serve in inpatient pediatrics hospital units [19].

Hein PT, *et al.* (2019) reported that; the rate of occupational violence among nurses was 72.7%. More than 75% of nurses witnessed violence, and the verbal assault was the most frequent. Around 25% of participants were victims of sexual assault. The regression study found that nurses working in the emergency room and outpatient clinic were 1.92 times more likely to have verbal harassment than people working in other units. They were also 3.02 times statistically more likely to have physical abuse [20].

Alkorashy, H. *et al.* (2016) reported that; about 50 % of participants had witnessed violence in the workplace during the 12 months before the study. Much of the respondents interpreted brutality in the workplace as physical harassment. Almost all nursing practitioners described patients as the leading cause. Slightly more than half of the understaffing, disagreements, long delays for service, and lack of staff preparation and crisis prevention policies have been listed as contributing factors [21].

Table 1. Author, country, year of publication, methodology, and results

Author, years, country	Methods and Objective	Outcomes
Gillespie, G. (2010) [16]	A qualitative study included 31 pediatric ED workers to describe the WPV that occurred in a pediatric ED and the negative effects on the workers.	WPV is an issue in pediatric ED and strategies are required to facilitate the wellbeing of staff and patients.
Li, Zhe, <i>et al.</i> (2017) China [17]	A retrospective cross-sectional study included 1,932 healthcare providers to evaluate the incidence, magnitude, consequences, and potential risk factors of workplace violence against the medical staff of children’s hospitals.	Pediatric staff is at high risk of violence. Hospital supervisors should pay attention to the consequences of WPV. There is a necessity for defensive measures to look after medical staff and provide a safer workplace environment.
Shaw J. (2015) [18]	A cross-sectional study involved 234 health care professionals explaining the ED staff’s perspectives on the possibility of occupational violence to direct changes and resolve employee complaints.	Workplace violence is growing in pediatric hospital emergency rooms. True and potential risks to the well-being of workers must be handled. The use of workers' expectations of risk and suggestions for change is a powerful method to direct the elimination of violence at work.
Strollo, Bonnie A., (2011) New York [19]	Qualitative research was used to describe nurses’ experience with workplace violence by directly examining their perceptions of the problem.	The findings of this research suggest that there is violence in the workplace and a great deal needs to be done to deter and control violence in the workplace toward nurses who serve in inpatient pediatrics hospital units.
Hein PT, <i>et al.</i> (2019) China [20]	The cross-section analysis involved 317 nurses in the period from January to December 2017 to measure the incidence of violence against nurses in the pediatric department and related factors.	The violence happened mainly in the emergency department and outpatient facilities. Verbal violations have been the most prominent.
Alkorashy, H. <i>et al.</i> (2016) Saudi Arabia [21]	A quantitative cross-sectional study on a sample of 370 nursing personnel to determine the prevalence rate of workplace violence against nursing professionals	The prevalence rate of WPV against nurses is extremely high among nurses in the studied sample

Violence against health care providers has been increasing in many countries. Exposure of pediatric staff to violence while carrying out their duties negatively affect nurses and physicians which may lead to loss of concentration while performing their duties and other negative consequences. Physicians are also a victim of occupational violence. Around a fifth of the emergency room physicians announced that they were victims of physical violence in the last year [22, 23]. Understanding the association between WPV and its effect on patient care is relevant because the results of the study could inspire hospital management to implement violence reduction initiatives. Gillespie's study findings indicate that all patients and visitors should be treated as though they have the potential to be violent. Employers must make sure the influence that violence has on employees' abilities to deliver patient care and debriefing and to learn how violence impacts patient care, how necessary it is to call for assistance when violence happens, and measures to avoid violence [16].

Several studies have reported high levels of WPV among health workers [24-29]. In a 2009 survey in Germany, 70.7 percent of healthcare professionals reported exposure to physical abuse, while 89.4% reported verbal violence, the repeated incidence of accidents and inadequate social assistance raised tension [30]. In Egypt, in the Ismailia Governorate report, the incidence of physical or verbal violence in nurses was 69.5% and 9.3% respectively [31]. The majority of the current study in China reports on tertiary and county-level hospitals emergency departments in general hospitals; research indicates that emergency departments have a high rate of WPV [32-35]. Several significant reports have suggested that nurses are at high risk of having WPV. The incidence rate of physical violence for nurses during the past 12 months, in Ethiopia, South Korea, Jordan, Germany ranged from 18.22 percent to 56.0 percent, the verbal abuse rate being from 63.8 percent to 89.58 percent, and the sexual harassment rate from 4.7 percent to 19.7 percent [36-41]. In a US report, approximately 25% of nurses serving in emergency departments were confronted with more than 20 physical violence incidents and 20% were confronted with more than 200 verbal violence incidents between 2006 and 2009. Many that were met with physical and/or verbal abuse also pointed to their fear of counterattack and lack of support for hospital management and ED management barriers to disclosing violence [42]. 88.1% of nurses reported verbal abuse from patients in a major mental health facility in Israel and 58.4% reported physical violence in the last year [43].

Research at Mansoura University Emergency Hospital has shown that only 7.4% of doctors did not experience abuse. Physical abuse was the most common with 76.5 percent followed by physical violence at 60.3 percent and sexual assault at 30.9 percent. Most physical and verbal aggressors are relatives/visitors to the patient. Hospital protection mechanisms are not available and there is no monitoring mechanism or survivor counseling for abuse of any sort [44]. In another survey performed in Ismailia in the emergency room, 59.7% of HCWs reported abuse. The bulk (58.2

percent) of verbal abuse was reported, while 15.7 percent reported physical violence. The investigators also found that inability to satisfy the needs of the patient and his families, and the amount of time they wait, are the key causes of abuse [45].

Study on the experience of nurses in these departments in the USA. Switzerland and Jordan have shown that they experience a greater rate of WPV than nurses in other units [46-48]. In a Palestinian government hospital, 80.4 percent of nurses reported having been subjected to abuse in the previous year; 20.8 percent were violent and 59.6 percent were non-physical [49]. A retrospective analysis in Australia found that verbal harassment (71%) was more common than physical abuse (29 percent) [50]. Research further indicates that 7.8% of nurses reported physically abusive experiences and 71.9% reported non-physically violent events in the previous year (a total of 588 nurses) [51]. Of the few population-based studies undertaken, a study of 1,404 health staff from Community Health Centers in Guangzhou and Shenzhen found that 51.64 percent witnessed WPV [52].

In previous research, 89% of violent cases were patients, 9% were family members, and 2% were patients' relatives [53]. In a nationwide survey conducted in the United States, 78% of emergency room physicians indicated that they had been the victim of occupational abuse in the previous year. Of these, 75 percent were in direct assaults, 21 percent were physical assaults, 5 percent were off-site confrontations, and 2 percent were abuse [54].

CONCLUSION

Pediatric physicians and nurses are exposed to different types of violence from parents or caregivers of the attending children in their departments due to the psychological stress caused by the condition of their child. Recommendations to teach pediatric staff how to deal with violent events and how to report them must be instructed clearly. Holding health education sessions to increase the public knowledge about the important and critical rules of pediatric staff as they deal with that delicate and vulnerable group of patients must be considered. Physicians and nurses also need to be well trained and acknowledged the anxiety and stress of parents about their children and how to break the bad news.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

1. United States Department of Labor. Safety and Health Topics | Workplace Violence | Occupational Safety and Health Administration [Internet]. [cited 2018 Sep 8]. Available from: <https://www.osha.gov/SLTC/workplaceviolence/>
2. Butchart A, Mikton C, Dahlberg LL, Krug EG. Global status report on violence prevention. 2014.
3. Algahtani FD. Healthy Lifestyle among Ha'il University Students, Saudi Arabia. *Int J Pharm Res Allied Sci.* 2020;9(1):160-7.

4. Hanawi SA, Saat NZ, Zulkafly M, Hazlenah H, Taibukahn NH, Yoganathan D, et al. Impact of a Healthy Lifestyle on the Psychological Well-being of University Students. *Int J Pharm Res Allied Sci.* 2020;9(2).
5. Sundus A, Ismail NE, Gnanasan S. Exploration of healthcare practitioner's perception regarding pharmacist's role in cancer palliative care, malaysia. *Pharmacophore.* 2018;9(4):1-7.
6. Shakeri H, Rahmanian V, Shakeri M, Mansoorian E. Study of anti-hbs antibody titer and associated factors among healthcare staff vaccinated against hepatitis B more than ten years in hospitals of Jahrom in 2016. *Pharmacophore.* 2018;9(1):156-61.
7. Phillips JP. Workplace violence against health care workers in the United States. *N Engl J Med.* 2016;374(17):1661-9.
8. McKoy Y, Smith MH. Legal considerations of workplace violence in healthcare environments. *In Nursing Forum 2001 (Vol. 36, No. 1, p. 5).* Blackwell Publishing Ltd.
9. Pinar R, Ucmak F. Verbal and physical violence in emergency departments: a survey of nurses in Istanbul, Turkey. *J Clin Nurs.* 2011;20(3-4):510-7.
10. Sisawo EJ, Ouédraogo SY, Huang SL. Workplace violence against nurses in the Gambia: mixed methods design. *BMC Health Serv Res.* 2017;17(1):311.
11. Lanctôt N, Guay S. The aftermath of workplace violence among healthcare workers: A systematic literature review of the consequences. *Aggress Violent Behav.* 2014;19(5):492-501.
12. Zafar W, Khan UR, Siddiqui SA, Jamali S, Razzak JA. Workplace violence and self-reported psychological health: coping with post-traumatic stress, mental distress, and burnout among physicians working in the emergency departments compared to other specialties in Pakistan. *J Emerg Med.* 2016;50(1):167-77.
13. Pan SJ, Cheng JM, Bai JG. Research on the causes of shortage of human resources in pediatrics and its countermeasures—a case study of Shanxi province. [J]. *Soft Sci Health.* 2016;30(7):31-4.
14. Pan DL. The dilemma of the children's medical resources shortage to be cracked [J]. *China Health Talent Resour.* 2013;9:17. (in Chinese).
15. Deng YQ, Ma XY, Cheng L. Study on workplace violence of pediatric nurses [J]. *Qilu J Nurs.* 2010;16(16): 51-3 (in Chinese).
16. Gillespie GL, Gates DM, Miller M, Howard PK. Violence against healthcare workers in a pediatric emergency department. *Adv Emerg Nurs J.* 2010;32(1):68-82.
17. Li Z, Yan CM, Shi L, Mu HT, Li X, Li AQ, et al. Workplace violence against medical staff of Chinese children's hospitals: a cross-sectional study. *PLoS One.* 2017;12(6):e0179373.
18. Shaw J. Staff perceptions of workplace violence in a pediatric emergency department. *Work.* 2015;51(1):39-49.
19. Strollo BA. Giving Voice: Pediatric Nurses' Perception of Workplace Violence. 2011.
20. Hein PT, Phuc DN, Trang LTT, Quang LN. Workplace Violence against Healthcare Nurses in the Vietnam National Children's Hospital: A Cross-Sectional Study. *J Health Hyg.* 2019;3(1):68-94.
21. Alkorashy HA, Al Moalad FB. Workplace violence against nursing staff in a Saudi university hospital. *Int Nurs Rev.* 2016;63(2):226-32.
22. Pompeii LA, Schoenfisch AL, Lipscomb HJ, Dement JM, Smith CD, Upadhyaya M. Physical assault, physical threat, and verbal abuse perpetrated against hospital workers by patients or visitors in six US hospitals. *Am J Ind Med.* 2015;58(11):1194-204.
23. Kowalenko T, Gates D, Gillespie GL, Succop P, Mentzel TK. Prospective study of violence against ED workers. *Am J Emerg Med.* 2013;31(1):197-205.
24. Chen ZH, Xia HY. Workplace violence against nurses in hospitals in Guangdong, China. *Inj Prev.* 2012;18(Suppl 1):A156-.
25. Talas MS, Kocaöz S, Akgüç S. A survey of violence against staff working in the emergency department in Ankara, Turkey. *Asian Nurs Res.* 2011;5(4):197-203.
26. Mroczek B, Mormul J, Kotwas A, Szkup M, Kurpas D. Patient aggression towards health care professionals. *Prog Health Sci.* 2014;4(2):67-74.
27. Raveesh BN, Lepping P, Lanka SV, Turner J, Krishna M. Patient and visitor violence towards staff on medical and psychiatric wards in India. *Asian J Psychiatr.* 2015;13:52-5.
28. Gomaa AE, Tapp LC, Luckhaupt SE, Vanoli K, Sarmiento RF, Raudabaugh WM, et al. Occupational traumatic injuries among workers in health care facilities—United States, 2012–2014. *MMWR Morb Mortal Wkly Rep.* 2015;64(15):405.
29. Lanza ML, Zeiss RA, Rierdan J. Non-physical violence: a risk factor for physical violence in health care settings. *AAOHN J.* 2006;54(9):397-402.
30. Franz S, Zeh A, Schablon A, Kuhnert S, Nienhaus A. Aggression and violence against health care workers in Germany—a cross sectional retrospective survey. *BMC Health Serv Res.* 2010;10(1):51.
31. Abbas MA, Fiala LA, Abdel Rahman AG, Fahim AE. Epidemiology of workplace violence against nursing staff in Ismailia Governorate, Egypt. *J Egypt Public Health Assoc.* 2010;85(1-2):29-43.
32. Liu H, Zhao S, Jiao M, Wang J, Peters DH, Qiao H, et al. Extent, nature, and risk factors of workplace violence in public tertiary hospitals in China: a cross-sectional survey. *Int J Environ Res Public Health.* 2015;12(6):6801-17.
33. Xing K, Jiao M, Ma H, Qiao H, Hao Y, Li Y, et al. Physical violence against general practitioners and nurses in Chinese township hospitals: a cross-sectional survey. *PLoS One.* 2015;10(11):e0142954.
34. Tingmei YA, Shuang LI, Guowei PA, Bo QU. Perceived Risk of Violence in Various Hospital Levels and De-partments in Urban and Rural China. *Iran J Public Health.* 2016;45(9):1238-40.
35. Chen WC, Hwu HG, Wang JD. Hospital staff responses to workplace violence in a psychiatric hospital in Taiwan. *Int J Occup Environ Health.* 2009;15(2):173-9.
36. Ferri P, Silvestri M, Artoni C, Di Lorenzo R. Workplace violence in different settings and among various health professionals in an Italian general hospital: a cross-sectional study. *Psychol Res Behav Manag.* 2016;9:263.
37. Fute M, Mengesha ZB, Waggari N, Tessema GA. High prevalence of workplace violence among nurses working at public health facilities in Southern Ethiopia. *BMC Nurs.* 2015;14(1):9.
38. Park M, Cho SH, Hong HJ. Prevalence and perpetrators of workplace violence by nursing unit and the relationship between violence and the perceived work environment. *J Nurs Scholarsh.* 2015;47(1):87-95.
39. Al-Omari H. Physical and verbal workplace violence against nurses in Jordan. *Int Nurs Rev.* 2015;62(1):111-8.
40. Schablon A, Zeh A, Wendeler D, Peters C, Wohlert C, Harling M, et al. Frequency and consequences of violence and aggression towards employees in the German healthcare and welfare system: a cross-sectional study. *BMJ Open.* 2012;2(5):e001420.
41. Hogarth KM, Beattie J, Morphet J. Nurses' attitudes towards the reporting of violence in the emergency department. *Australas Emerg Nurs J.* 2016;19(2):75-81.
42. Gacki-Smith J, Juarez AM, Boyett L, Homeyer C, Robinson L, MacLean SL. Violence against nurses working in US emergency departments. *JONA: J Nurs Admin.* 2009;39(7/8):340-9.
43. Itzhaki M, Peles-Bortz A, Kostitsky H, Barnoy D, Filshtinsky V, Bluvstein I. Exposure of mental health nurses to violence associated with job stress, life satisfaction, staff resilience, and post-traumatic growth. *Int J Ment Health Nurs.* 2015;24(5):403-12.
44. Elkhawaga G, El-Gilany AH, Abd-El-Raouf SE. Violence against physicians in Mansoura University Emergency Hospital, Egypt. *Egypt J Community Med.* 2012;30(3).
45. Abdellah RF, Salama KM. Prevalence and risk factors of workplace violence against health care workers in emergency department in Ismailia, Egypt. *Pan Afr Med J.* 2017;26(1):1-8.
46. Pompeii LA, Schoenfisch AL, Lipscomb HJ, Dement JM, Smith CD, Upadhyaya M. Physical assault, physical threat, and verbal abuse perpetrated against hospital workers by patients or visitors in six US hospitals. *Am J Ind Med.* 2015;58(11):1194-204.
47. Hahn S, Müller M, Hantikainen V, Kok G, Dassen T, Halfens RJ. Risk factors associated with patient and visitor violence in general hospitals: Results of a multiple regression analysis. *Int J Nurs Stud.* 2013;50(3):374-85.
48. ALBashtawy M, Aljezawi ME. Emergency nurses' perspective of workplace violence in Jordanian hospitals: A national survey. *Int Emerg Nurs.* 2016;24:61-5.
49. Kitaneh M, Hamdan M. Workplace violence against physicians and nurses in Palestinian public hospitals: a cross-sectional study. *BMC Health Serv Res.* 2012;12(1):1-9.
50. Cashmore AW, Indig D, Hampton SE, Hegney DG, Jalaludin BB. Workplace violence in a large correctional health service in New

- South Wales, Australia: a retrospective review of incident management records. *BMC Health Serv Res.* 2012;12(1):245.
51. Jiao M, Ning N, Li Y, Gao L, Cui Y, Sun H, et al. Workplace violence against nurses in Chinese hospitals: a cross-sectional survey. *BMJ Open.* 2015;5(3).
 52. Lin WQ, Wu J, Yuan LX, Zhang SC, Jing MJ, Zhang HS, et al. Workplace violence and job performance among community healthcare Workers in China: the mediator role of quality of life. *Int J Environ Res Public Health.* 2015;12(11):14872-86.
 53. Kowalenko T, Walters BL, Khare RK, Compton S, Michigan College of Emergency Physicians Workplace Violence Task Force. Workplace violence: a survey of emergency physicians in the state of Michigan. *Ann Emerg Med.* 2005;46(2):142-7.
 54. Behnam M, Tillotson RD, Davis SM, Hobbs GR. Violence in the emergency department: a national survey of emergency medicine residents and attending physicians. *J Emerg Med.* 2011;40(5):565-79.