# Investigating Endurance Training with Ergometer Bicycle in Changes the Plasma Levels of Chemerin on Overweight Women

Ahmad Reza Askari Ashtiani<sup>1</sup>\*, Rezvane Galdavi<sup>2</sup>, Mehri Jafari<sup>3</sup>

<sup>1</sup>Health Promotion Research Center of Zahedan Medical Sciences University, Zahedan, Iran. <sup>2</sup>Department of Sport Sciences, University of Birjand, Birjand, Iran. <sup>3</sup>Department of Physical Education and Sport Sciences, University of Sistan and Baluchestan, Zahedan, Iran.

#### Abstract

Adipokine chemerin is a pre-inflammation associated with obesity and metabolic syndrome. The present study aimed to investigate the changes in the plasma levels of chemerin of women with overweight and obese during a period of training on a cycle-ergometer. This study was quasi-experimental type and was performed on 22 overweight and obese women (BMI>25Kg/m<sup>2</sup>) who were within the age range of 30-45 in Zahedan. Subjects were selected targeted and using permuted block randomization was divided into two groups of subjects including training groups and control groups. The training protocol that was performed lasted for 8 weeks (3 sessions per week) by cycling on a cycle-ergometer (60-75% heart rate reserve) with a duration of 16 to 30 minutes. Blood samples were collected in two stages before and after the same conditions to measure plasma levels of chemerin. To analyze the results, t-test, ANOVA, covariance, and Kruskal-Wallis analysis were used at a significant level of  $\alpha < 0.05$ . Intra-group comparisons showed that 8 weeks of aerobic exercise with ergometer bicycle in addition to a significant reduction of obesity-related factors significantly reduced plasma levels of chemerin in training groups (p=0.002). A significant difference between the changes in plasma levels of chemerin of the studied groups was observed during the inter-group comparison. To reduce the pre-inflammatory factor of chemerin and obesity-related factors in overweight and obese women, training on the cycle-ergometer has been considered as a preventive approach.

Keywords: Chemerin, Obesity and Overweight, Cycle-ergometer

#### INTRODUCTION

Obesity and overweight are now a major concern worldwide [1] because obesity is reported to be the cause of diseases such as type 2 diabetes, coronary artery disease, cardiovascular disease [2], and different cancers [3]. Therefore, the development of prevention programs and fighting obesity is of particular importance. Researchers link obesity with lifestyle variables such as physical activity patterns [4] and factors such as disturbed nutritional behaviors, stress, etc. [5]. Today, adipose tissue, as an active endocrine system, by producing fatty acids and secreting several types of proteins regulates hormones of absorbing and consuming energy. Some of the hormones secreted from adipose tissue, which play an important role in the process of inflammation and cardiovascular disease, are called adipokine [6]. In obese people, the production of most adipokines that contribute to the balance and regulation of energy, immunity, metabolism and fatty homeostasis and angiogenesis [7] has increased or impaired, and all of the above factors increase the risk of cardiovascular disease [7] and disorders such as type 2 diabetes, inflammation, and metabolic syndrome [8]. Chemerin is an adipokine that is initially secreted as an immature and inactive peptide with 163 amino acids and a molecular weight of 18 KiloDalton (kDa) from white and liver adipose tissue in the form of proxamethrin. Then it turns into mature and active chemerin with 143 amino acids and a molecular weight of 16 kDa by extracellular serine protease enzyme and is found in plasma and serum [9]. Chemerin has localized effects on adipogenesis and potentially profound effects on metabolism and inflammation [10]. Due to its role in fat synthesis, energy metabolism, inflammation, and the invocation of immune cells to the site of injury and inflammation have been of great importance [11]. Production of chemerin is associated with the volume of adipose tissue, so that the higher the adipose tissue, the, and this higher chemerin secretion at lipogenesis level is associated with insulin resistance [12]. Based on the

Address for correspondence: Ahmad Reza Askari Ashtiani, Health Promotion Research Center of Zahedan Medical Sciences University, Zahedan, Iran. Ahmadaskary@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**How to cite this article:** Ashtiani ARA, Galdavi R, Jafari M. Investigating Endurance Training with Ergometer Bicycle in Changes the Plasma Levels of Chemerin on Overweight Women. Arch Pharm Pract. 2021;12(1):50-4. https://doi.org/10.51847/y6IUtBmY4U results of recent studies, the serum level of chemerin in patients with obesity and type 2 diabetes increases. In addition, the amount of chemerin has a positive correlation with BMI, waist circumference, blood pressure, triglyceride, LDL-C, and a negative correlation with HDL-C and adiponectin [13]. As physical inactivity is an important factor in the incidence of obesity [14], one of the methods treatments of obesity has heavily focused on sports activities [15]. This is because exercise affects the amount of adipose tissue that affects adipocytic secretions [16] and also, reducing fat percent decreases levels of inflammation [17]. Given the importance and potential role of sports exercises in reducing weight and complications of overweight, improving cardiovascular [18] and oxidative stress reduction [19] are used today to improve the physical condition of obese and diabetic patients, instead of using chemical drugs- regular exercise, including endurance exercises.

Nowadays, exercise on a cycle-ergometer is considered as a new exercise among the people of the community. However, the effect of this new exercise on the concentration of adipokine chemerin and the relationship between this type of exercise and the mechanism of this hormone is not clear due to a lack of research history. Several studies have been conducted on the effects of various types of sports activities on chemerin levels and the results have not been consistent due to differences in the type of subjects, intensity, type, and volume of exercise. Zolfaghari et al. [20] and Askari et al. [21] reported no significant changes in chemerin levels of their subjects, but Jafari et al. [22] and Sadeghipour et al. [23] reported a significant decrease in chemerin levels in their studies. Therefore, considering the contradictions in the results of the effect of exercise on chemerin levels and the gap of studies in the effect of training on cycle-ergometer on the plasma levels of chemerin, conducting this study seems necessary.

# MATERIALS AND METHODS

The present study is a randomized clinical trial (RCT) (laboratory) using pre-test and post-test stages along with experimental and control groups. At first, with the call of research at the level of the offices of Zahedan, overweight and obese women volunteer to participate were selected in a targeted way. At a briefing session, some explanations were given on the objectives of the plan and how it was going to be performed, then the questions and ambiguities were answered, and the subjects completed the consent form with full knowledge and completed the questionnaire containing personal information, medical-health, physical activity, etc. The inclusion criteria for entering the research were women aged 30-45 years, having a general physical and mental health level BMI>25 (Kg/m<sup>2</sup>), lack of participation in regular exercise activities during the past 6 months, lack of cardiovascular disease, diabetes, hormonal disorders, kidney and liver diseases, surgery, tobacco use and any therapeutic intervention affecting laboratory results. Exclusion criteria included not following the diet, participation in other sports

activities, illness during exercise, pregnancy, and irregular attendance of subjects in one of the groups tested. To assess physical and mental health, the participants were examined by the physician and accepted in case of proven health. Subjects were selected targeted and using permuted block randomization were divided into two groups of 11 subjects including training group 1 (endurance exercise with cycleergometer) and group2 (control group). Then, the individual and physical characteristics of the subjects including age. height in cm (without shoes, legs stuck together, and the buttocks, shoulders, and back in contact with Height gauge), weight in kilograms (light clothes without shoes), BMI by dividing body weight in kilograms by squared height in meters, waist circumference in the tibia, hip circumference in the most prominent part of the pelvis by the strip meter, WHR from splitting waist circumference to hip and body fat percentage by measuring subcutaneous fat using a calipers model (SAEHAN Made in Korea) at three points, triceps, above the pelvis and the thigh on the right side of the body were measured and for calculating body fat percentage were given to Jackson and Pollack formula [24].

All measurements were done one stage before the beginning of the exercise (24 hours before), and the next step after the end of the training (the day after the last training session) in the morning and the same conditions. To assess the changes in plasma levels of chemerin, blood samples were taken in the pre-test (24 hours before the beginning of the first session) and post-test (48 hours after the last training session to prevent the effect of acute inflammation induced by exercise) in the same conditions after 12 hours of fasting [25]. To prevent the effects of night-daytime oscillations on plasma levels of Chemerin, sampling was performed at a specific time of day (7 to 8 in the morning) at different stages [26]. Blood samples were centrifuged for 5 minutes at a speed of 3000 RPM and stored frozen at -20°C for plasma separation. Eastbiopharm human kit made by a Chinese-American company was used for biochemical analysis and measurement of the plasma levels of chemerin with a sensitivity of 4.99 ng/L. Two weeks before the start of the training, all subjects were asked to take no medication during the study period and refrain from participating in other training programs, and observe the usual diet during the study period. The protocol was run for 8 weeks and 3 sessions per week. The training program included 10 minutes of warm-up running at low intensity and tensile strength at the beginning of the session, followed by a specific training protocol and the end of each training session, cooling to 10 minutes by soft running and stretching movements. Specific exercise protocols included cycling on a cycle-ergometer (Sport Cross-Life Care model) with a reserved heart rate of 60 to 75% [27]. The Pulse Oximeter A310 fingerprint speedometer made in Germany was used to control the intensity of heart rate during exercise. The intensity of exercise was calculated based on the reserved heart rate through the Kornen formula [28].

Heart rate during exercise= (maximum heart rate - resting heart rate)  $\times$  intended heart rate (60% to 75%) + Rest heart rate

The maximum heart rate was calculated from the formula (220-age). The duration of the main training at the start of the course was 16 minutes, which gradually increased to 30 minutes by the end of the training period. In each week of the training, based on the principle of overload, the intensity and duration of the training (2 minutes per week and 5% every two weeks) increased [27].

#### Statistical Method

Descriptive statistics were used to categorize raw data, to determine the mean and standard deviations, and to determine the normality of the data related to the subjects of the research groups. Shapiro-Wilk test was used. To study the intra-group differences, a paired t-test was used, and for inter-group comparison, one-way analysis of variance (ANOVA), covariance, Kruskal-Wallis, and if significant, Tukey's post hoc test was used to determine differences. SPSS 16 was used

to perform statistical calculations and the statistical significance level was considered 0.05.

## **RESULTS AND DISCUSSION**

The mean and standard deviation of subjects' characteristics in the pretest phase are presented in **Table 1**. One-way ANOVA showed that there was no significant difference in the basic levels of personal data (P<0.05), but the mean of chemerin before the intervention was significantly different between the groups (P = 0.025).

Table 1. Descrip	tive Indices	of Individual							
Characteristics in the Two Study Groups									
Group	(1)	(2)							
Variable	M±SD	M±SD							
Age(Year)	34.90±5.97	36.00±6.24							
Height(centimeter)	160.83±3.60	160.27±4.73							
Weight(Kilograms)	$83.49{\pm}10.88$	77.25±7.89							
BMI (Kg / m2)	31.71±5.08	30.70±3.79							

Table 2. The mean and standard deviati	on (±) of the variables of the second s	he research according to	dependent t-test,
one-way analysis of variance, covariance	, and Kruskal-Wallis		

Variable G	roups	Pre-test	Post-test	Intra-group P	Inter-group P
Weight	(1)	81.22±12.76	76.90±11.57	*0.001	V 0 007
(Kilograms)	(2)	77.25±7.89	76/79±8.22	0.332	¥ 0.997
Body fat percentage	(1)	41.59±3.89	37.37±3.72	*0.001	V. 0. (70)
	(2)	39.61±3.34	39.12±3.37	0.066	¥ 0.679
BMI	(1)	31.46±5.08	29.81±4.67	*0.001	
( Kg/m <sup>2</sup> )	(2)	30.20±2.58	30.01±2.70	0.319	¥ 0.910
WHR	(1)	$0.78{\pm}0.05$	0.77±0.05	0.111	# 0.486
	(2)	$0.81 \pm 0.08$	$0.80 \pm 0.08$	0.242	
chemerin	(1)	330.36±54.20	293.86±48.47	*0.001	** • • • • •
(ng/L)	(2)	341.43±61.81	341.71±57.55	0.870	~\$<0.001
Values are shown as mean $\pm$ standard deviation	* Signif	icant statistical significance	(P <0.05) \$ Covaria	nce test # Kruskal W	allis Test ¥ One-

Values are shown as mean  $\pm$  standard deviation \* Significant statistical significance (P < 0.05) \$ Covariance test # Kruskal Wallis Test ¥ Or way analysis of variance

**Table 2** shows that intra-group changes using paired t-test significantly reduced in three groups (1), in obesity-related factors such as weight, body fat percentage, and BMI. Plasma levels of chemerin decreased in the group (1), which were statistically significant. (P<0.05). In group 2, weight, fat mass, BMI, and chemerin plasma levels did not show significant differences (P>0.05). Based on the covariance test, there was a significant difference between the changes in the Chemerin plasma levels in the studied groups (p<0.05). However, the results of ANOVA showed no significant differences between the changes from pre to post-test of other anthropometric variables in the studied groups (P>0.05).

## CONCLUSION

Because of the lack of a research background on the effect of training on cycle-ergometer on plasma levels of chemerin, in the present discussion, we have used studies related to the subject in some way. Also, in this study, the effect of endurance training on cycle-ergometer on plasma levels of chemerin on overweight and obese women was examined from two different perspectives. With this explanation, of the most important findings of the present study, we can refer to the significant reduction of plasma levels of Chemerin in the group (1) following 8 weeks of training on cycle-ergometer with 60% to 75% reserved heart rate. Other significant results were a significant reduction in weight, fat percentage, and

BMI in the group (1) at the end of the period compared to the base values. However, WHR values did not change significantly in any of the groups compared to baseline values. Adipose tissue plays an important role in the development of metabolic problems associated with obesity with the production of inflammatory factors [29]. As stated, obesity is one of the factors leading to a rise in chemerin levels in humans, and a decrease in the weight is due to lower levels of serum levels of chemerin [30]. Although the correlation of chemerin with fat accumulation is not known well, the results of the studies show that chemerin concentration is associated with a high-fat percentage. WHR [31], BMI, and waist circumference [13]. Shin et al. examined the relationship between the levels of chemerin and body composition in 173 women, and found it in correlation with BMI, lipid percentage, WHR, visceral fat, fasting insulin, total cholesterol, and triglyceride, so introduced chemerin as an indicator associated with visceral obesity in cardiovascular disease [32]. Other studies also show that active lifestyle [33], weight loss due to caloric restriction, obstructive surgery [10], and regular exercise [34] leads to a reduction of serum chemerin. These researchers also attributed a significant decrease in chemerin levels in overweight and obese subjects to lower body fat levels and weight loss following exercise. By exercising, skeletal muscle capacity increases for using fats, which can play an important role in controlling the weight of obese and overweight people and reducing cardiovascular risk factors. It can be justified that a significant proportion of the fatty acids required in the muscles in operation is provided by increasing the lipid triglyceride lipolysis of the adipose tissue 3 to 4 times. Sports activity doubles the amount of blood flow to the adipose tissue and increases the amount of blood to the active muscle of the body. This reduces body fat and improves body composition by eliminating the balance between intake and consumption of energy and creating a negative calorie balance that itself may lead to a decrease in chemerin of plasma after exercise [35].

Another possible mechanism for decreasing chemerin can be the increase in caloric expenditure after training and its effect on reducing the pathway of adipogenesis. Since with an increase in lipogenesis, chemerin values decrease, and this decrease reflects the reduction in the rate of adipogenicity, which is an important indicator for diabetic and cardiovascular patients. Various studies have suggested the origin of chemerin to be adipose tissue, which can affect atherosclerosis development as paracrine and may result in the call of macrophages and inflammatory responses in atherosclerotic plaques [36]. Saremi et al. (2010) have considered a significant reduction in plasma levels of chemerin following endurance training related to weight loss, especially visceral fat in obese individuals. They believe that changes in visceral fat after exercise may have an important role in regulating the secretion of macrophages into fat tissue and inflammatory markers such as chemerin [18]. On the other hand, the results of this study conflict with the research

by Ghanbarzadeh *et al.* [27], Asgari *et al.* [34], and Moradi *et al.* [37].

In the research by Ganbarzadeh et al., which was performed on elderly women (HRmax intensity of 61-88% in the period of 30 to 16 minutes), despite a significant decrease in weight and BMI, there was no significant change in the plasma levels of chemerin, fat percentage and WHR of the subjects [29]. The reason for the difference in the results of the research mentioned in this study can be at the level of fitness, age, the health status of the subjects, and other factors. As in the study by Ghanbarzadeh et al. the age range of subjects was 55-70, while the age range of the subjects in this study was 30-45. Likely, the differences in age and physiological conditions of the subjects in the mentioned studies have had a different effect on the secretion and metabolic systems of these subjects. It seems that the response of the chemerin hormone to exercise is different subjects is different due to age. In the study by Asgari et al. [34], the subjects were young girls with an average age of 20 years. The reasons for the inconsistency of the results with this study can be the factorial difference between the subjects' age and differences in the practice protocol. In the study by Moradi et al. [37], which was performed on inactive elderly men, there was no significant change in factors such as weight, fat percentage, BMI, and consequently in the plasma levels of chemerin. According to the results of various studies, which have linked the reduction of chemerin to reduction of fat mass, and as Alfadda et al. showed a positive correlation between the serum levels of chemerin in adult women and men with different degrees of obesity with BMI [38], lack of changes in serum levels of chemerin in the subjects of the study by Moradi et al. can be attributed to weight and fat mass remained constant. Among the limitations of the present study, we can mention the low volume of samples, stressors, endocrine discharges, the genetic characteristics of the subjects, and the inability to control the diet of the subjects accurately by the researchers.

## CONCLUSION

It seems that doing eight weeks of endurance training on cycle-ergometer can be considered as a new and effective method for decreasing plasma levels of chemerin and improving the body composition in obese and overweight women. To reach the definitive result, given the lack of study in this area, it is suggested that the effect of this exercise should be examined in other humans and obese and overweight men.

ACKNOWLEDGMENTS: The present study was the result of a research project approved by the Health Promotion Research Center of Zahedan University of Medical Sciences with a clinical practice record number IRCT2016020826443N1, which was co-sponsored by the deputy of research in Zahedan University of Medical Sciences. We would like to express our appreciation and gratitude to this center and the vice president of research at the university. Moreover, we sincerely appreciate the assistance and cooperation of the subjects and all those who helped us in the implementation of this research project. CONFLICT OF INTEREST: None FINANCIAL SUPPORT: None ETHICS STATEMENT: None

## References

- Oqden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. JAMA. 2006;295(13):1549-55.
- Marinou K, Tousoulis D, Antonopoulos AS, Stefanadi E, Stefanadis C. Obesity and cardiovascular disease: from pathophysiology to risk stratification. Int J Cardiol. 2010;138(1):3-8.
- Rose DP, Vona-Davis L. Interaction between menopausal status and obesity in affecting breast cancer risk. Maturitas. 2010;66(1):33-8.
- Singh GK, Kogan MD, Van Dyck PC, Siahpush M. Racial/ethnic, socioeconomic, and behavioural determinants of childhood and adolescent obesity in the United States: analyzing independent and joint associations. Ann Epidemiol. 2008;18(9):682-95.
- 5. Stunkard AJ. Depression and obesity. Biol Psychiatry. 2003;54(3):330-7.
- Vendrell J, Broch M, Vilarrasa N, Molina A, Gómez JM, Gutiérrez C, et al. Resistin, adiponectin, ghrelin, leptin, and proinflammatory cytokines: relationships in obesity. Obes Res. 2004;12(6):962-71.
- 7. Ronti T, Lupattelli G, Mannarino E. The endocrine function of adipose tissue: an update. Clin Endocrinol. 2006;64(4):355-65.
- Gustafson B. Adipose tissue, inflammation and atherosclerosis. J Atheroscler Thromb. 2010;17(4):332-41.
- 9. Bozaoglu K, Bolton K, McMillan J, Zimmet P, Jowett J, Collier G, et al. Chemerin is a novel adipokine associated with obesity and metabolic syndrome. Endocrinology. 2007;148(10):4687-94.
- Chakaroun R, Raschpichler M, Klöting N, Oberbach A, Flehmig G, Kern M, et al. Effects of weight loss and exercise on chemerin serum concentrations and adipose tissue expression in human obesity. Metabolism. 2012;61(5):706-14.
- 11. Nehal Hamdy ES, Wadie Fawzy M, Nagwa AEG, Ezzat A. Role of serum chemrin level in atherosclerosis in egyptian patients with type 2 diyabetes mellitues. J Appl Sci Res. 2013;9:775-8.
- Sell H, Laurencikiene J, Taube A, Eckardt K, Cramer A, Horrighs A, et al. Chemerin is a novel adipocyte-derived factor inducing insulin resistance in primary human skeletal muscle cells. Diabetes. 2009;58(12):2731-40.
- 13. Saremi A, Moslehabadi MF, Parastesh M. Effects of Twelve-week strength training on serum chemerin, TNF— $\alpha$  and CRP level in subjects with the metabolic syndrome. Iran J Endocrinol Metab. 2011;12(5):536-43.
- Watts K, Jones TW, Davis EA, Green D. Exercise training in obese children and adolescents: Current Concepts. Sports Med. 2005;35(5):375-92.
- Hassink SG, Zapalla F, Falini L, Datto G. Exercise and the obese child. Prog Pediatr Cardiol, 2008;25(2):153-7.
- Chtara M, Chamari K, Chaouachi M, Chaouachi A, Koubaa D, Feki Y, et al. Effects of intra-session concurrent endurance and strength training sequence on aerobic performance and capacity. Br J Sports Med. 2005;39(8):555-60.
- Hosseini Kakhak A, Amiri Parsa T, Haghighi A, Asgari R, Chamri M, Hedayati M. The effect of resistance training on hs- CRP and cystatin C in obese adolescents. Bimonthly Scientific-Research Shahed University. 2009;85.
- Saremi A, Shavandi N, Parastesh M, Daneshmand H. Twelve-week aerobic training decreases chemerin level and improves cardiometabolic risk factors in overweight and obese men. Asian J Sports Med. 2010;1(3):151.
- Hajer GR, Van Haeften TW, Visseren F. Adipose tissue dysfunction in obesity, diabetes, and vascular diseases. Eur Heart J. 2008;29(24):2959-71.

- Zolfaghari M, Taghian F, Hedayati M. The effects of green tea extract consumption, aerobic exercise and a combination of these on chemerin levels and insulin resistance in obese women. Iran J Endocrinol Metab. 2013;15(3):253-61.
- Asgari R, Ravasi A, Gaeini A, Hedyati M, Hamedinia M. The effect of combined exercise training on indices adipokines and insulin sensitivity in overweight women. Sport Biom Sci. 2011;1(5):25-34.
- Jafari M, Mogharnasi M, Salimi Khorashad A. Effects of Endurance and Resistance Training on Plasma Levels of Chemerin and Factors Related to Obesity in Overweight and Obese Females. Armaghanedanesh. 2015;20(4): 273-86.
- 23. Sadeghipour HR, Daryanoosh F, Salesi M. The Effect of 12 Weeks of Aerobic Interval Training on Chemerin and Vaspin Serum Concentrations and Insulin Resistance Index in Overweight Students. J Zanjan Univ Med Sci Health Serv. 2015;23(96):78-88.
- Jackson AS, Pollock ML, Ward AN. Generalized equations for predicting body density of women. Med Sci Sports Exerc. 1980;12(3):175-81.
- Jahangard T, Torkaman G, Ghoosheh B, Hedayati M, Dibaj A. The effect of short-term aerobic training on coagulation and fibrinolytic factors in sedentary healthy postmenopausal women. Maturitas. 2009;64(4):223-7.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clin Chem. 1972;18(6):499-502.
- Ghanbarzadeh M, Kazemi AR. The Comparison of Three Different Concurrent Training on Chemerin Plasma Levels, Insulin Resistance and Physical Performance in Elderly Women. J Knowl Health. 2016;10(4):40-7.
- Saghebjoo M, Dastigerdi S, Afzalpour ME, Hedayati M. Effects of aerobic and resistance training on plasma visfatin levels in overweight women. Koomesh. 2012;13(2):225-32.
- Maury E, Brichard S. Adipokine dysregulation, adipose tissue inflammation and metabolic syndrome. Mol Cell Endocrinol 2010;314(1):1-16.
- 30. Becker M, Rabe K, Lebherz C, Zugwurst J, Göke B, Parhofer KG, et al. Expression of human chemerin induces insulin resistance in the skeletal muscle but does not affect weight, lipid levels, and atherosclerosis in LDL receptor knockout mice on high-fat diet. Diabetes. 2010;59(11):2898-903.
- Ernst MC, Sinal CJ. Chemerin: at the crossroad of inflammation and obesity. Trends Endocrinol Metab. 2010;21(11):660-7.
- 32. Shin HY, Lee DC, Chu SH, Jeon JY, Lee MK, Im JA, et al. Chemerin levels are positively correlated with abdominal visceral fat accumulation. Clin Endocrinol. 2012;77(1):47-50.
- Neuparth MJ, Proença JB, Santos-Silva A, Coimbra S. The positive effect of moderate walking exercise on chemerin levels in Portuguese patients with type 2 diabetes mellitus. J Investig Med. 2014;62(2):350-3.
- Asgari R, Ravasi AA, Gaeini AA, Hedayati M, Hamedinia MR. The effect of combined and endurance exercise training on some adipokines, GH and lipid profile in overweight females. J Sport Biosci. 2015;6(4):399-413.
- Yamaner F, Bayraktaroğlu T, Atmaca H, Ziyagil MA, Tamer K. Serum leptin, lipoprotein levels, and glucose homeostasis, between national wrestlers and sedentary males. Turk J Med Sci. 2010;40(3):471-7.
- Goralski KB, McCarthy TC, Hanniman EA, Zabel BA, Butcher EC, Parlee SD, et al. Chemerin, a novel adipokine that regulates adipogenesis and adipocyte metabolism. J Biol Chem. 2007;282(38):28175-88.
- 37. Moradi F, Heydarzadeh A, Vosoughi-Baneh V. The effect of an endurance training program on serum levels of leptin and chemerin adipokines in inactive lean men. Feyz. 2014;18(5):419-27.
- Alfadda AA, Sallam RM, Chishti MA, Moustafa AS, Fatma S, Alomaim WS, et al. Differential patterns of serum concentration and adipose tissue expression of chemerin in obesity: adipose depot specificity and gender dimorphism. Mol Cells. 2012;33(6):591-6.