

# The Effect of Nitrous Oxide on Postoperative Pain Third Molar Surgery

Nasser Kaviani<sup>1</sup>, Ablofath Hosseinzadeh<sup>2</sup>, Ali Akhavan<sup>3\*</sup>, Mozafar Mehraban<sup>4</sup>

<sup>1</sup>Anesthesiologist- Assistant professor, Department of Oral and Maxillofacial Surgery and Dental Research Center, School of Dentistry- Isfahan University of Medical Sciences, Isfahan, Iran. <sup>2</sup> Assistant Professor. Department of Oral and Maxillofacial Surgery, and Torabinejad Dental Research Center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran. <sup>3</sup> Dental Materials Research Center, Department of Endodontics, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran. <sup>4</sup> Graduate, Dental Student Research Committee, School of dentistry, Isfahan University of Medical Sciences, Isfahan, Iran.

## Abstract

**Introduction:** The postoperative pain is one of the most important factors influencing fear and anxiety. The use of inhalation sedation makes a patient relax and pain-free during surgery. This study aimed to assess the effects of 50% nitrous oxide on postoperative pain after third molar teeth surgery. **Materials and Method:** Forty-three adult patients were involved in this prospective randomized controlled clinical trial. All patients were of ASA1 status. The subjects were divided randomly into two groups receiving 100% oxygen or 50% nitrous oxide/50% oxygen. The degree of the postoperative pain was assessed at different periods: 6 hours after the surgery, 12 hours after surgery, and 24 hours after surgery by VAS. The number of oral analgesic (Ibuprofen 400mg) taken during the first 24 hours was also assessed in the two groups. Data were analyzed with Chi-Square and Independent Sample T-test. **Results:** The pain score and the amount of total analgesic consumption in the nitrous oxide groups were significantly less than the oxygen group ( $P < 0.005$  respectively). The mean number of oral analgesics taken by the patients in the nitrous oxide and oxygen groups was 5.27 and 7.9 respectively. **Conclusion:** The admiration of nitrous oxide/oxygen mixture during third molar surgery can decrease postoperative pain.

**Keywords:** Dentistry, Nitrous Oxide, Pain, Third Molar

## INTRODUCTION

Worldwide the number of surgical operations to remove wisdom teeth is immense, in England alone, approximately 63,000 are removed in National Health Service (NHS) hospitals each year. Many patients need time off work and their quality of life is significantly affected. However, despite these consequences, people are often most concerned about pain following the operation, which can be severe. It was suggested that the most intense pain is felt three to five hours after surgery. The pain experienced after oral surgery is widely used as a model to measure the effectiveness of painkillers in general<sup>[1]</sup>.

Oral surgery procedures are often associated with fear and anxiety<sup>[2-4]</sup>. Postoperative pain is one of the most important factors influence dental fear. Pain and anxiety control during oral surgery can decrease postoperative pain<sup>[5]</sup>. Age, gender, smoking, duration of surgery, type and amount of local anesthesia and severity of tissue damage during the surgery are responsible for the postoperative pain. Many pharmacological and non-pharmacological methods are applied for postoperative pain control. The non-steroid anti-inflammatory drugs (NSAIDs) are the most common medications prescribed<sup>[5, 6]</sup>. NSAIDs are highly effective and have not sedative effect. The adverse effects of NSAIDs

are gastric ulceration, renal impairment, and coagulopathy. Nitrous oxide (N<sub>2</sub>O) is widely used as an anesthetic gas and as an inhalational sedation agent for day-stay minor surgical procedures in the various medical disciplines. It has good analgesic properties. The analgesic effect of 50% N<sub>2</sub>O is equal to 10 milligrams of intramuscular morphine<sup>[7]</sup>. Inhalation sedation with N<sub>2</sub>O/O<sub>2</sub> is one of the main techniques used for relief of pain and anxiety during minor dental procedures<sup>[5, 8-10]</sup>. It has a rapid onset, short, and provides for excellent control of the pain, anxiety, and

**Address for correspondence:** Ali Akhavan, Dental Materials Research Center, Department of Endodontics, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran.  
E-mail: akhavan@dnt.mui.ac.ir

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work noncommercially, as long as the author is credited and the new creations are licensed under the identical terms.

**How to cite this article:** Kaviani, N., Hosseinzadeh, A., Akhavan, A., Mehraban, M., The Effect of Nitrous Oxide on Postoperative Pain Third Molar Surgery. Arch Pharma Pract 2018;9(2):5-8.

awareness during the uncomfortable dental procedures<sup>[11-13]</sup>. The main disadvantage of inhalation sedation with N<sub>2</sub>O/O<sub>2</sub> is that many patients are not adequately sedated by it. In an animal study, Goto<sup>[14]</sup> showed that 30% and 75% N<sub>2</sub>O administered before a noxious stimulation produced a preemptive analgesic effect (decrease postoperative pain). Also, Ong<sup>[15]</sup> showed that 50% N<sub>2</sub>O administration before dental surgery has no effect on postoperative pain in humans. Venchard<sup>[16]</sup> showed that sedation for dental surgery was improved by combining nitrous oxide and intravenous midazolam and found this technique to be safe and reliable, requiring reduced doses of midazolam and it made improvement in patient recovery and cooperation.

This study aimed to evaluate the effect of 50% N<sub>2</sub>O administration on postoperative pain and the need for analgesic in third molar teeth surgery.

## MATERIALS AND METHODS

After the approval of research in the research deputy of Isfahan Dental school, (No: 386002). In this prospective randomized controlled clinical trial, Forty-three patients from patients who referred to the department of oral and maxillofacial surgery of Isfahan University of Medical Sciences for removing their impacted third molar teeth. The Inclusion criteria were age between 20 to 40 years, no history of cardiovascular and pulmonary disease, and have not a contraindication of N<sub>2</sub>O administration. After taking a complete medical and dental history and informed consent, the patients were randomly divided into two groups, based on a random digit table: the control group who received 100% O<sub>2</sub> and the study group who received 50% N<sub>2</sub>O / 50% O<sub>2</sub> through a nasal mask during procedure.

All patients were of ASA1 status. Exclusion criteria for the study were personality or psychiatric disorders, nasal abnormalities (septal deviations, inferior conchal hypertrophy) chronic obstructive pulmonary disease, and ASA 2 through 5 statuses. The same surgeon performed all the surgical procedures while an anesthesiologist controlled the sedation. Heart rate, respiratory rates, and oxygen saturation were monitored for both groups during surgery. After five minutes of N<sub>2</sub>O gas administration, the local anesthetic (lidocaine 2% + epinephrine 1/100,000) was injected into the operative site. After surgery, the patient point VAS for intraoperative pain, the postoperative pain intensity was assessed by using the visual analog scale (VAS) system at different periods: 6 hours after the surgery, 12 hours after surgery, and 24 hours after surgery. The patient indicated pain intensity on a line of 10 cm long (0 no pain to 10 worst pain). The amount of the analgesic (Ibuprofen 400 mg) consumed during the first 24 hours was also recorded.

## RESULTS

The Forty- three patients who were involved in the study were divided into two groups: 100% oxygen and 50% N<sub>2</sub>O/ 50% O<sub>2</sub>.

The study group included 22 patients (11 male and 11 female). With a mean age of 26.6±5.5 years. The control group included 21 patients (8 male and 13 female) with a mean age of 26.5±6.1 years (Table: 1). Besides, in each group, 13 patients had a history of dental treatment and the others had not. There was no significant difference in age, gender, and history of dental treatment between two groups (Chi-Square, P = 0.85). Therefore, the influence of age, gender, and history of dental treatment had little effect on the treatment outcomes.

The mean number of local anesthetic cartilage used in the control and study groups was 2.57 and 2.05 respectively. The mean time of surgery was 33.1 for the study group and 32.3 for the control group and not different in two groups (p>0.05). At the end of the surgery, the patients were asked to provide an overall evaluation of the pain experience during surgery on a line of 10 cm long (0 = no pain and 10 =worst pain possible)

The intensity of pain during surgery, the intensity of pain during the first 24 hours after surgery, and amount of analgesic consumption during the first 24 hours after the surgery was significantly higher in the control group compared to the study group (Independent Sample T-test P<0.000, P < 0.009 and P <0.005 respectively). (Tables: 2, 3 and 4).

The sedation score during surgery was higher in the study group. (Mann Whitney, P=0.000). (Table: 5)

The patient had not any medical complications such as hypoxia, hypertension, and hypotension during treatment.

## DISCUSSION

Many animals and humans studies showed that local anesthesia could provide preemptive analgesia by preventing central sensitization. Kissin<sup>[17]</sup> stated that the preemptive analgesia block would need to be of sufficient strength and duration to be effective. Another study by Gorden<sup>[18]</sup> demonstrated that blockade of postoperative pain during the first 4 hours with bupivacaine following oral surgery resulted in less postoperative pain 1 to 2 days after surgery. Concerning nitrous oxide, there is paucity in researches that study the preemptive analgesic effect of N<sub>2</sub>O in humans. Venchard<sup>[16]</sup> found that combining nitrous oxide and Midazolam improved patient's sedation during oral surgery and reduced doses of Midazolam administered to them. Patients were initially titrated with 10% nitrous oxide, increasing by increments of 10% up to a maximum of 40% nitrous oxide and 60% oxygen. Midazolam was then titrated (initially 2 mg wait 2 min with increments of 1 mg every minute until appropriately sedated) whilst still administering

40% nitrous oxide. Goto<sup>[14]</sup> has shown that N<sub>2</sub>O produces a dose-dependent preemptive analgesic effect in rats. In his study, he showed that 20 minutes of 30% or 75% nitrous oxide administered before a noxious stimulation (formalin injection), produced a lasting effect on the pain behavior in the animal. A recent study by Ong<sup>[15]</sup> showed that 50% N<sub>2</sub>O administered preoperatively for 20 minutes does not have any preemptive analgesic effect in patients having impacted third molar teeth extraction under local anesthesia. In our study, 50% of nitrous oxide was used to produce conscious sedation. Our results are consistent with the results of the study of Goto<sup>[14]</sup> and Venchard<sup>[16]</sup> but contradict these of Ong<sup>[15]</sup>. In Ong<sup>[15]</sup> study, nitrous oxide was given for 20 minutes before the surgical procedure. In our study, the nitrous oxide was started 5 minutes before the local anesthesia administration and continued throughout the surgical procedure. The mean time of the surgical procedures was 32.7 minutes.

The administration of N<sub>2</sub>O for a longer time (more than 20 minutes) may have led to the difference in results between our study and the Ong<sup>[15]</sup> study. As was stated by Kissin<sup>[17]</sup>, sufficient strength and duration are needed for the analgesic block to be effective. Extending the N<sub>2</sub>O administration into the intraoperative period may have provided adequate time for blocking the development of central sensitization. The intraoperative pain and anxiety may also influence the postoperative pain. By using N<sub>2</sub>O for patient's sedation during the surgical procedures, the patient and the surgeon are more comfortable and the patients may experience less intraoperative pain. We have also found that the patient's intraoperative pain in the study group was significantly lower than that of the control group ( $P < 0.005$ ). So, we can also conclude that N<sub>2</sub>O sedation may have decreased the postoperative pain by improving the conditions of the surgical procedure for the patient and the surgeon. As stated before, due to the lack of adequate studies on the preemptive analgesic effect of N<sub>2</sub>O in humans, further assessment and larger studies are necessary to confirm our findings.

## CONCLUSION

The result of our study showed that using 50% N<sub>2</sub>O administered five minutes before surgery for removal of impacted teeth and continued until the end of the operation, could significantly decrease postoperative pain and the number of analgesics taken by patients.

## Acknowledgements

We thank the research deputy of Isfahan Dental School for cooperation. Also, for Dr. Hosseinzadeh, who died in a traumatic accident, we wish peace.

## REFERENCES

1. Bailey E, Worthington H, Coulthard P. Ibuprofen and/or paracetamol (acetaminophen) for pain relief after surgical removal of lower wisdom teeth, a Cochrane systematic review. *Br Dent J* 2014;216(8):451-5.
2. Kyle, B. N., D. W. McNeil, B. Weaver, and T. Wilson. "Recall of dental pain and anxiety in a cohort of oral surgery patients." *Journal of dental research* 95, no. 6 (2016): 629-634.
3. Santuchi, Camila Carvalho, Sheila Cavalca Cortelli, José Roberto Cortelli, Luís Otávio Miranda Cota, Camila Oliveira Alencar, and Fernando Oliveira Costa. "Pre - and post - treatment experiences of fear, anxiety, and pain among chronic periodontitis patients treated by scaling and root planing per quadrant versus one - stage full - mouth disinfection: a 6 - month randomized controlled clinical trial." *Journal of clinical periodontology* 42, no. 11 (2015): 1024-1031.
4. Tarazona, Beatriz, Pablo Tarazona-Álvarez, David Peñarrocha-Oltra, Juan Rojo-Moreno, and María A. Peñarrocha-Diago. "Anxiety before extraction of impacted lower third molars." *Medicina oral, patología oral y cirugía bucal* 20, no. 2 (2015): e246.
5. Vassend, Olav. "Anxiety, pain and discomfort associated with dental treatment." *Behaviour research and therapy* 31, no. 7 (1993): 659-666.
6. Prause G. 15 years of preoperative ambulatory care in Graz. *Anaesthetist* 1994; 43(4): 223-8.
7. Parbrook, G. D., G. A. D. Rees, and G. S. Robertson. "Relief of post-operative pain: Comparison of a 25% nitrous-oxide and oxygen mixture with morphine." *British medical journal* 2, no. 5407 (1964): 480.
8. Kaufman, E., D. C. Chastain, A. M. Gaughan, and R. H. Gracely. "Staircase assessment of the magnitude and time-course of 50% nitrous-oxide analgesia." *Journal of dental research* 71, no. 9 (1992): 1598-1603.
9. Veerkamp, J. S., R. J. Gruythuisen, and J. Hoogstraten. "Anxiety reduction with nitrous oxide: a permanent solution?." *ASDC journal of dentistry for children* 62, no. 1 (1995): 44-48.
10. Trapp, Larry D. "Mechanisms of acute pain: an update." *CDA. JOURNAL* 34, no. 12 (2006).
11. Clark, M., and A. Brunick. "Nitrous Oxide Interaction with the Body." Clark M, Brunick A-Nitrous Oxide and Oxygen Sedation. St. Louis, Mosby (1999): 103-114.
12. LUHMANN, JAND, ROBERT M. KENNEDY, David M. Jaffe, and JOHN D. McALLISTER. "Continuous-flow delivery of nitrous oxide and oxygen: a safe and cost-effective technique for inhalation analgesia and sedation of pediatric patients." *Vacuum* 3 (1999): 30LPM.
13. Annequin, Daniel, Ricardo Carbajal, Pierre Chauvin, Olivier Gall, Barbara Tourniaire, and Isabelle Murat. "Fixed 50% nitrous oxide oxygen mixture for painful procedures: a French survey." *Pediatrics* 105, no. 4 (2000): e47-e47.
14. Goto, Takahisa, J. J. Marota, and Gregory Crosby. "Nitrous oxide induces preemptive analgesia in the rat that is antagonized by halothane." *Anesthesiology* 80, no. 2 (1994): 409-416.
15. Ong, K. S., R. A. Seymour, and J. M. L. Tan. "A prospective randomized crossover study of the preemptive analgesic effect of nitrous oxide in oral surgery." *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology* 98, no. 6 (2004): 637-642.
16. Venchard, G. R., P. J. Thomson, and R. Boys. "Improved sedation for oral surgery by combining nitrous oxide and intravenous Midazolam: a randomized, controlled trial." *International journal of oral and maxillofacial surgery* 35, no. 6 (2006): 522-527.
17. Kissin, Igor. "Preemptive analgesia." *Anesthesiology: The Journal of the American Society of Anesthesiologists* 93, no. 4 (2000): 1138-1143.
18. Gordon, Sharon M., Jaime S. Brahim, Ronald Dubner, Linda M. McCullagh, Christine Sang, and Raymond A. Dionne. "Attenuation of pain in a randomized trial by suppression of peripheral nociceptive activity in the immediate postoperative period." *Anesthesia & Analgesia* 95, no. 5 (2002): 1351-1357.

**Table 1:** Characteristics of two volunteer groups (M=male, F=female)

Group	Gender M/F	Age
50% N <sub>2</sub> O	11/11	26.6(±5.5)
100% O <sub>2</sub>	8/13	26.5(±6.1)
<b>P Value</b>	0.85	0.85

**Table 2:** The amount of pain intensity during surgery (VAS= Visual Analog Scale).

	Control group	Study group	p value
<b>Pain</b>	2.88(±1.09)	4.88(±1.13)	0.000

**Table 3:** The amount of pain intensity in 2 groups (VAS=Visual Analog Scale ; h=hours)

	Control group	Study group	p value
<b>Pain during first 6h</b>	4.09(±0.89)	3.35(±0.8)	<0.009
<b>Pain during first 12h</b>	3(±0.75)	1.95(±0.8)	<0.009
<b>Pain during first 24h</b>	1.6(±0.66)	0.97(±25)	<0.009

**Table 4:** Amount of analgesic consumption in the first 24 hours

	Control group (number of tablets)	Study group (number of tablets)	p- Value
<b>First 6h</b>	2.86(±1.01)	1.77(±0.60)	<0.005
<b>First 12h</b>	5.21(±1.09)	3.23(±0.75)	<0.005
<b>First 24h</b>	7.9(±1.86)	5.27(±1.03)	<0.005

**Table 5:** Sedation score in tow groups

	Control group (number of patients)	Study group (number of patients)	p- Value
<b>Fully awake</b>	20	2	0.000
<b>Drowsy</b>	1	4	0.000
<b>Eye closed responded to verbal command</b>	0	8	0.000
<b>Eye closed responded to mild physical stimulus</b>	0	8	0.000