

Third-Generation Cephalosporin-Resistant Enterobacteriaceae in Africa

Dear Editor,

Recent studies by Zhang et al.^[1] and Grundman et al.^[2] added to the evidence regarding the global dissemination of antibiotic resistance among Enterobacteriaceae species in China and Europe, respectively. The African continent is not left out, because it also belongs to the worldwide trend of growing antibiotic resistance among Enterobacteriaceae such as *Escherichia coli* and *Klebsiella* spp., particularly to third-generation cephalosporin (3GC).^[3] 3GC-resistant Enterobacteriaceae are responsible for a range of infections, which include recurrent urinary tract infections, complicated device-associated infections, and fatal bloodstream infections in the hospital and community. These infections, subsequently, lead to prolonged therapy, the consumption of limited healthcare resources, and increased mortality. In recent years, the high levels of 3GC-resistant Enterobacteriaceae have been reported in African countries, which are mediated by the expression of extended-spectrum beta-lactamases ESBL and plasmid AmpC beta-lactamases. About 62.15% of ESBL-producing Enterobacteriaceae isolated from a hospital in Mthatha, South Africa,^[4] exhibited resistance to 3GCs. A similar trend was also reported in Zimbabwe (24.5%),^[5] Algeria (69.85%),^[6] and Togo (95.62%).^[7]

The emergence and spread of 3GC-resistant Enterobacteriaceae in African countries is worrisome because these pathogens often exhibit coresistance to the other classes of antibiotics. This further restricts physicians' treatment options and reduces the effectiveness of 3GCs in the future. Moreover, the next available treatment options, which are the newer antibiotics such as carbapenems, are very costly and not readily available in most countries in Africa. The emergence of resistance to these antibiotics of last resort in certain parts of Africa also worsens the already critical situation. One of the major contributing factors to the rising trend of 3GC-resistant Enterobacteriaceae in African countries is the misuse and overuse of this class of antibiotics for the empirical treatment of bacterial infections and for postsurgical prophylaxis. The misuse is further enhanced by their unregulated sales and widespread consumption without a physician's prescription. Other factors include a lack of antimicrobial surveillance strategies and limited human and healthcare resources.

The World Health Organisation's global priority list of antibiotic-resistant bacteria to guide research, discovery, and the development of new antibiotics published in February 2017 identified a critical need for the development of new antibiotics for Enterobacteriaceae resistant to 3GC.^[3] In addition, drastic intervention and multidisciplinary strategies need to be established in many regions and countries in Africa to prevent and tackle infections caused by antibiotic-resistant pathogens. These include prioritizing infection prevention and control in hospitals, the development and implementation of a strong regulatory policy on antibiotic sales and prescription, and lastly a strengthening of regional and national antimicrobial surveillance.

Acknowledgments

I thank Tahir Mehmood Khan, Syed Tabish Razi Zaidi, Rahul P. Patel, and Long Chiau Ming for their comments and suggestions.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Ifunanya E. Asogwa

Faculty of Pharmacy, University of Benin, Benin City, Nigeria

Address for correspondence: Dr. Ifunanya Evangeline Asogwa, Faculty of Pharmacy, University of Benin, Benin City, Nigeria. E-mail: linann3000@yahoo.com

REFERENCES

1. Zhang Y, Jin L, Ouyang P, Wang Q, Wang R, Wang J, Gao H, Wang X, Wang H. Evolution of hypervirulence in carbapenem-resistant *Klebsiella pneumoniae* in China: a multicentre, molecular epidemiological analysis. *Journal of Antimicrobial Chemotherapy*. 2019 Nov 12.
2. Grundmann H, Glasner C, Albiger B, Aanensen DM, Tomlinson CT, Andrasević AT, Cantón R, Carmeli Y, Friedrich AW, Giske CG, Glupczynski Y. Occurrence of carbapenemase-producing *Klebsiella pneumoniae* and *Escherichia coli* in the European survey of carbapenemase-producing Enterobacteriaceae (EuSCAPE): a prospective, multinational study. *The Lancet Infectious Diseases*. 2017 Feb 1;17(2):153-63.
3. World Health Organisation. Global Priority List of Antibiotic Resistant Bacteria to Guide Research, Discovery and Development of New Antibiotics; 2017. Available from: <http://www.who.int/medicines/>

- publications/WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf?ua=1. [Last accessed on 03 Mar 2017].
4. Vasaikar S, Obi L, Morobe I, Bisi-Johnson M. Molecular characteristics and antibiotic resistance profiles of Klebsiella isolates in Mthatha, Eastern Cape Province, South Africa. *International journal of microbiology*. 2017;2017.
 5. Magwenzi MT, Gudza-Mugabe M, Mujuru HA, Dangarembizi-Bwakura M, Robertson V, Aiken AM. Carriage of antibiotic-resistant Enterobacteriaceae in hospitalised children in tertiary hospitals in Harare, Zimbabwe. *Antimicrobial Resistance & Infection Control*. 2017 Dec;6(1):10.
 6. Lagha N, Hassaine H, Robin F, Bonnet R, Abdelouahid D. Prevalence and molecular typing of extended-spectrum b-lactamases in Escherichia coli, Enterobacter cloacae and Citrobacter freundii isolates from Laghouat Hospital, Algeria. *Afr J Microbiol Res* 2016;10:1430-8.
 7. Salah FD, Diagbouga S, Dabire AM, Sadjji AY, Nadembega C, Moumouni A, et al. First detection of resistance genes encoding extended spectrum beta-lactamase producing Escherichia coli at Lome, Togo. *Arch Clin Microbiol* 2016;7. doi: 10.4172/1989-8436.100062.

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work noncommercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to cite this article: Asogwa IE. Third-Generation Cephalosporin-Resistant Enterobacteriaceae in Africa. *Arch Pharma Pract* 2018; 9(1):1-2.

© 2018 Archives of Pharmacy Practice